

Hodge theory and deformations of SKT manifolds

Gil R. Cavalcanti*
Department of Mathematics
Utrecht University

Abstract

We use tools from generalized complex geometry to develop the theory of SKT (a.k.a. pluriclosed Hermitian) manifolds and more generally manifolds with special holonomy with respect to a metric connection with closed skew-symmetric torsion. We develop Hodge theory on such manifolds showing how the reduction of the holonomy group causes a decomposition of the twisted cohomology. For SKT manifolds this decomposition is accompanied by an identity between different Laplacian operators and forces the collapse of a spectral sequence at the first page. Further we study the deformation theory of SKT structures, identifying the space where the obstructions live. We illustrate our theory with examples based on Calabi–Eckmann manifolds, instantons, Hopf surfaces and Lie groups.

Contents

1	Linear algebra	3
2	Intrinsic torsion of generalized Hermitian structures	8
2.1	The Nijenhuis tensor	9
2.2	The intrinsic torsion and the road to integrability	10
2.3	The operators δ_{\pm} and $\bar{\delta}_{\pm}$	12
3	Parallel Hermitian and bi-Hermitian structures	12
4	SKT structures	14
5	Hodge theory	19
5.1	Differential operators, their adjoints and Laplacians	19
5.2	Signature and Euler characteristic of almost Hermitian manifolds	20
5.3	Hodge theory on parallel Hermitian manifolds	22
5.4	Hodge theory on SKT manifolds	23
5.5	Relation to Dolbeault cohomology	23
5.6	Hermitian symplectic structures	27
6	Hodge theory beyond $U(n)$	27
6.1	Integrability	29

Keywords. Strong KT structure, generalized complex geometry, generalized Kähler geometry, Hodge theory, instantons, deformations.

2010 *Mathematics subject classification.* 53C55; 53C15; 53C29; 53D30.

*gil.cavalcanti@gmail.com

7	Instantons over complex surfaces	29
8	Stability	32
8.1	Stability of SKT structures I	32
8.2	Stability of generalized Kähler structures	38
8.3	Stability of SKT structures II	40
9	Appendix - SKT differentials	41

Introduction

Looking beyond the Levi–Civita connection in Riemannian geometry, one finds a number of other metric connections with interesting properties. Normally these families of connections are defined by some characteristic of the torsion tensor and recurrent themes of research are connections with parallel torsion or connections with skew symmetric torsion. The “strong” torsion condition refers to the latter: a *strong torsion* connection on a Riemannian manifold is a metric connection whose torsion is skew symmetric and closed. In this setting, a *Kähler structure with strong torsion*, or *SKT structure* is a Hermitian structure (g, I) together with a strong torsion connection for which I is parallel. A weaker notion is that of a (strong) parallel Hermitian structure which for us means a connection with closed, skew symmetric torsion for which the holonomy is $U(n)$. The difference between a parallel Hermitian structure and an SKT structure being that in the former case integrability of the complex structure is not required.

A reason to study of such objects comes from string theory, where closed 3-forms arise naturally as fields in their sigma models [28, 33]. Once a 3-form is added to the sigma model, if one still requires a nontrivial amount of supersymmetry, the type of geometry of the target space has to move away from the usual Kähler geometry. It was precisely following this path that Gates, Hull and Roček [15] discovered the bi-Hermitian geometry that nowadays also goes by the name of generalized Kähler geometry [20] as the solutions to the $(2, 2)$ -supersymmetric sigma model. Requiring less supersymmetry without giving up on the idea altogether leads one to consider models where there is more left than right supersymmetry or models where the right side is simply absent. These conditions lead to $(2, 1)$ or $(2, 0)$ supersymmetric sigma models and supersymmetry holds if and only if the target space has an SKT structure [24]. This point of view also leads one to consider parallel Hermitian and bi-Hermitian structures as these are geometric structures imposed by a sigma model with an extended supersymmetry algebra [10, 11].

Mathematical properties of SKT structures have been subject of study of several papers since the 90’s, including those of Bismut [3], Grantcharov *et al* [18], Fino and collaborators [12, 13] and more recently Streets and Tian [30, 31]. Yet, until now there was no framework in which basic Kähler results, such as Hodge theory and deformation theory, obtained meaningful counterparts in the SKT world. Indeed the opposite seemed to be the case: Since SKT manifolds are, in particular, complex, their space of forms inherits a natural bi-grading, but a simple check in concrete examples shows that there is no corresponding decomposition of their cohomology. Further, by studying these structures on six dimensional nilmanifolds Fino, Parton and Salamon [12] produced examples showing that their Frölicher spectral sequence does not necessarily degenerate at the first page, they do not satisfy the dd^c -lemma, manifolds carrying these structures may not be formal and that these structures are not stable under deformations of the complex structure. In short, several Kähler properties seem to have been lost once the torsion was included. Put another way, given a complex manifold, we were left with no tools to decide whether it admitted an SKT metric or not.

Here we tackle SKT structures from a new point of view. The key observation is that SKT structures have yet another description, this time, as a ‘generalized structure’, i.e., a geometric structure on $TM = TM \oplus T^*M$. In fact, in this paper we show that, in a very precise way, SKT structures lie half way between generalized Hermitian and generalized Kähler structures. Using this

approach, we show that some of the negative results mentioned earlier have a positive counterpart involving the torsion while those with negative answers obtain a conceptual explanation for their failure to hold.

Indeed, the first observation is that, as structures defined on $\mathbb{T}M$ with the H -Courant bracket, the natural differential operator to consider is $d^H = d + H\wedge$, where $H = d^c\omega$, and ω is the Hermitian form. Hence questions about d , and its decomposition as $\partial + \bar{\partial}$ miss an important ingredient and were doomed from the start. The cohomology of d^H is only \mathbb{Z}_2 -graded, yet we show that a parallel Hermitian structure induces a $\mathbb{Z} \times \mathbb{Z}_2$ grading on the space of forms which itself induces a $\mathbb{Z} \times \mathbb{Z}_2$ grading on the d^H -cohomology. This is achieved by introducing the *intrinsic torsion* of a generalized almost Hermitian structure and using it to show that the d^H -Laplacian preserves a $\mathbb{Z} \times \mathbb{Z}_2$ -graded decomposition of the space of forms. For SKT manifolds one can go further and prove an identity of Laplacians, extending Gualtieri's work on generalized Kähler geometry [19]. This way we relate a cohomology naturally defined in terms of the SKT data with the d^H -cohomology. These identities imply further that a variation of the Frölicher spectral sequence degenerates at the first page. As an application, we return to the moduli space of instantons on a bundle over a compact complex surface and show that the existence of an SKT structure in this space (obtained by Lübke and Teleman [27] by an *ad hoc* method) can be seen as a consequence of the Hodge theory developed for SKT manifolds.

With this new framework at hand, we study of the problem of stability of SKT structures. Precisely, we use the operators introduced by the generalized complex setup to determine the obstruction space to the stability problem. In its classical version, we prove that if the Dolbeault cohomology group $H^{1,2}(M)$ vanishes, then any deformation of the complex structure can be accompanied by a deformation of the metric so that the pair remains SKT.

This paper is organized as follows. In Section 1 we develop the linear algebra pertinent to generalized complex, generalized Hermitian and SKT structures. In particular we show that an SKT structure gives rise to a $\mathbb{Z} \times \mathbb{Z}_2$ -grading on the space of forms. In Section 2 we introduce the intrinsic torsion of a generalized Hermitian structure and, in Sections 3 and 4, we relate SKT structures and parallel Hermitian structures to the vanishing of certain components of the intrinsic torsion. In Section 5 we study Hodge theory for SKT and parallel Hermitian structures and prove that in both cases the d^H -cohomology decomposes according to the decomposition of forms induced by the structure. As an application, in Section 7 we recover the result that the moduli space of instantons over a complex surface has an SKT structure. In Section 8 we use our framework to study the problem of deformations of SKT and generalized Kähler structures.

Acknowledgements: This research was supported by a Marie Curie Intra European Fellowship and a VIDI grant from NWO, the Dutch science foundation. The author is thankful to Anna Fino, Sönke Rollenske, Ulf Lindstrom, Martin Roček, Stefan Vandoren and Maxim Zabzine for useful conversations.

1 Linear algebra

Given a vector space V^m we let $\mathbb{V} = V \oplus V^*$ be its “double”. \mathbb{V} is endowed with a natural symmetric pairing:

$$\langle X + \xi, Y + \eta \rangle = \frac{1}{2}(\eta(X) + \xi(Y)), \quad X, Y \in V \quad \xi, \eta \in V^*.$$

Elements of \mathbb{V} act on $\wedge^\bullet V^*$ via

$$(X + \xi) \cdot \varphi = i_X \varphi + \xi \wedge \varphi.$$

One can easily check that for $v \in \mathbb{V}$

$$v \cdot (v \cdot \varphi) = \langle v, v \rangle \varphi,$$

hence $\wedge^\bullet V^*$ is naturally a module for the Clifford algebra of \mathbb{V} . In fact, it is the space of spinors for $Spin(\mathbb{V})$ and hence comes equipped with a spin invariant pairing, the *Chevalley pairing*:

$$(\varphi, \psi)_{Ch} = -(\varphi \wedge \psi^t)_{top},$$

where \cdot^t indicates transposition, an \mathbb{R} -linear operator defined on decomposable forms by

$$(\alpha_1 \wedge \cdots \wedge \alpha_k)^t = \alpha_k \wedge \cdots \wedge \alpha_1,$$

and *top* means taking the degree m component.

The spin group, $Spin(\mathbb{V})$, acts on both \mathbb{V} and on spinors in a compatible manner, namely, its action on \mathbb{V} is by conjugation using Clifford multiplication

$$g_* v = g v g^{-1} \quad \text{for all } g \in Spin(\mathbb{V}), v \in \mathbb{V}$$

and on $\wedge^\bullet V^*$ by the Clifford action described above, so we have

$$g \cdot (v \cdot \varphi) = (g v g^{-1}) \cdot (g \varphi) = (g_* v) \cdot (g \cdot \varphi) \quad (1.1)$$

for all $g \in Spin(\mathbb{V})$, $v \in \mathbb{V}$ and $\varphi \in \wedge^\bullet V^*$.

Example 1.1 (*B-field transform*). Given $B \in \wedge^2 V^* \subset \mathfrak{spin}(\mathbb{V})$, it acts on \mathbb{V} . Namely, we can regard B as a map from V into V^*

$$B(X + \xi) = -i_X B = -B(X).$$

Exponentiating this map we have

$$e_*^B : \mathbb{V} \longrightarrow \mathbb{V} \quad e_*^B(X + \xi) = X + \xi - B(X).$$

And this action is compatible with the action of $e^B \in Spin(\mathbb{V})$ on forms:

$$e^B : \varphi \longrightarrow e^B \wedge \varphi.$$

So we have $e_*^B : \wedge^k \mathbb{V} \longrightarrow \wedge^k \mathbb{V}$ and for $\alpha \in \wedge^\bullet \mathbb{V}$ and $\varphi \in \wedge^\bullet V^*$

$$(e_*^B \alpha) \cdot e^B \varphi = e^B (\alpha \cdot \varphi).$$

■

We will be interested in introducing geometric structures on \mathbb{V} . The first we consider is a generalized metric, as introduced by Gualtieri [20].

Definition 1.2. A *generalized metric* on V is an automorphism $\mathcal{G} : \mathbb{V} \longrightarrow \mathbb{V}$ which is orthogonal and self-adjoint with respect to the natural pairing and for which the bilinear tensor

$$\langle \mathcal{G}v, w \rangle, \quad v, w \in \mathbb{V}$$

is positive definite.

Since \mathcal{G} is orthogonal and self-adjoint, we have $\mathcal{G}^{-1} = \mathcal{G}^t = \mathcal{G}$, hence $\mathcal{G}^2 = \text{Id}$. Therefore \mathcal{G} splits \mathbb{V} into its ± 1 -eigenspaces: $\mathbb{V} = V_+ \oplus V_-$ and the projection $\pi_V : \mathbb{V} \longrightarrow V$ gives isomorphisms $\pi : V_\pm \longrightarrow V$. Further, given a generalized metric \mathcal{G} we can write $\mathbb{V} = \mathcal{G}V^* \oplus V^*$ and $\mathcal{G}V^*$ is isomorphic to V via the projection $\pi_V : \mathbb{V} \longrightarrow V$. Since both V and $\mathcal{G}V$ are isotropic subspaces of \mathbb{V} which project isomorphically onto V , we can describe $\mathcal{G}V$ as the graph of a linear map $b : V \longrightarrow V^*$, that is $b \in \otimes^2 V^*$. Isotropy means that $b \in \wedge^2 V^*$ and hence gives rise to an orthogonal transformation

of the natural pairing, e_*^b . This map has the property that $e_*^b : \mathcal{G}V^* \rightarrow V$, hence, after an orthogonal transformation of \mathbb{V} , we can assume that $\mathcal{G}V^* = V$. For this splitting,

$$\mathcal{G} = \begin{pmatrix} 0 & g \\ g^{-1} & 0 \end{pmatrix} \quad (1.2)$$

where g is an ordinary metric on V . The splitting of \mathbb{V} determined by a generalized metric is the *metric splitting*.

If V is endowed with an orientation, we can define a generalized *Hodge star operator* [19] as follows. Since $\pi_V : V_+ \rightarrow V$ is an isomorphism, V_+ inherits an orientation. Then we let $\{e_1, e_2, \dots, e_m\}$ be a positive orthonormal basis of V_+ and let $\star = -e_m \cdots e_2 \cdot e_1 \in \text{Clif}(\mathbb{V})$. Then

$$\star\varphi := \star \cdot \varphi,$$

where \cdot denotes Clifford action.

With this definition, we have

$$(\varphi, \star\varphi)_{Ch} > 0 \quad \text{if } \varphi \neq 0 \quad (1.3)$$

If the splitting of \mathbb{V} is the metric splitting, we have

$$(\varphi, \star\psi)_{Ch} = \varphi \wedge \star\psi,$$

where $*$ is the usual Hodge star, hence, in this splitting, \star is the usual Hodge star except for a change in signs given by the Chevalley pairing. Since $\star = -e_m \cdots e_2 \cdot e_1$, we have that

$$\star^2 = (-1)^{\frac{m(m-1)}{2}}$$

and hence it splits the space of forms into its eigenspaces, namely, into its ± 1 -eigenspaces $\wedge_{\pm}^{\bullet} V^*$ if m is zero or one modulo four or its $\pm i$ -eigenspaces if n is 2 or 3 modulo 4. This allows us to define *self-dual* and *anti self-dual forms* in all dimensions:

Definition 1.3. We say that a form $\varphi \in \wedge^{\bullet} V_{\mathbb{C}}^*$ is SD if $\star\varphi = -i^{\frac{m(m-1)}{2}}\varphi$ and that it is ASD if $\star\varphi = i^{\frac{m(m-1)}{2}}\varphi$. We denote the space of SD-forms by $\wedge_{+}^{\bullet} V^*$ and the space of ASD-forms by $\wedge_{-}^{\bullet} V^*$.¹

The Clifford action of elements in V_{\pm} either preserves or switches the eigenspaces of \star :

Lemma 1.4. Let $v_{\pm} \in V_{\pm}$. Then acting via Clifford action on forms we have

$$v_+ \star = (-1)^{m-1} \star v_+ \quad \text{and} \quad v_- \star = (-1)^m \star v_-$$

hence for m even,

$$V_+ : \wedge_{\pm}^{\bullet} V^* \rightarrow \wedge_{\mp}^{\bullet} V^* \quad \text{and} \quad V_- : \wedge_{\pm}^{\bullet} V^* \rightarrow \wedge_{\pm}^{\bullet} V^*$$

and for m odd

$$V_+ : \wedge_{\pm}^{\bullet} V^* \rightarrow \wedge_{\pm}^{\bullet} V^* \quad \text{and} \quad V_- : \wedge_{\pm}^{\bullet} V^* \rightarrow \wedge_{\mp}^{\bullet} V^*$$

Proof. For $v_- \in V_-$, we have that $\langle v_-, V_+ \rangle = 0$ hence v_- graded commutes with \star . For $v_+ \in V_+$, we can choose an orthonormal basis for V_+ for which $e_1 = v_+ / \|v_+\|$, so that v_+ anti commutes with all the remaining elements of the basis and commutes with e_1 . \square

For the rest of this section we will introduce structures on V which force its dimension to be even so we let $m = 2n$.

¹The choice of signs on the Chevalley pairing and of \star were made so that on a four manifold, the notion of SD and ASD agrees with the usual notions of self-dual and anti self-dual on 2-forms.

Definition 1.5. A *generalized complex structure* on V is a complex structure on \mathbb{V} which is orthogonal with respect to the natural pairing. A *generalized Hermitian structure* or a $U(n) \times U(n)$ *structure* on V is a generalized complex structure \mathcal{J}_1 on V and a generalized metric \mathcal{G} such that \mathcal{J}_1 and \mathcal{G} commute.

Given a generalized complex structure \mathcal{J} on V , we can split $\mathbb{V}_{\mathbb{C}}$, the complexification of \mathbb{V} , into the $\pm i$ -eigenspaces of \mathcal{J} : $\mathbb{V}_{\mathbb{C}} = L \oplus \bar{L}$. The spaces L and \bar{L} are maximal isotropic subspaces of $\mathbb{V}_{\mathbb{C}}$ such that $L \cap \bar{L} = \{0\}$. Since the natural pairing is nondegenerate, we can use it to identify $\bar{L} = L^*$. Precisely, we identify

$$v \mapsto 2\langle v, \cdot \rangle \in L^*, \quad \text{for } v \in \bar{L}. \quad (1.4)$$

Given a generalized Hermitian structure $(\mathcal{J}_1, \mathcal{G})$ on V , $\mathcal{J}_2 = \mathcal{G}\mathcal{J}_1$ is orthogonal with respect to the natural pairing and squares to $-\text{Id}$, hence it is also a generalized complex structure. Since $\pi_V : V_{\pm} \rightarrow V$ are isomorphisms, and $\mathcal{J}_1|_{V_{\pm}}$ is a complex structure on V_{\pm} orthogonal with respect to the natural pairing, it induces complex structures I_{\pm} on V compatible with the metric g induced by \mathcal{G} making V into a bi-Hermitian vector space. We can further form the corresponding Hermitian forms $\omega_{\pm} = g \circ I_{\pm}$.

Given any generalized Hermitian structure, $\wedge^{\bullet} V_{\mathbb{C}}^*$ splits as the intersections of the eigenspaces of \mathcal{J}_1 and \mathcal{J}_2 : $U^{p,q} = U_{\mathcal{J}_1}^p \cap U_{\mathcal{J}_2}^q$, where $U_{\mathcal{J}_i}^p$ is the ip -eigenspace of \mathcal{J}_i in $\wedge^{\bullet} V_{\mathbb{C}}^*$. In this context, the generalized Hodge star is related to the action of $\mathbb{J}_i = e^{\frac{\pi \mathcal{J}_i}{2}}$, namely:

Lemma 1.6. (Gualtieri [19]) In a generalized Hermitian vector space one has

$$\star = -\mathbb{J}_1 \mathbb{J}_2.$$

This means that we can read the decomposition of forms into SD and ASD from the $U^{p,q}$ decomposition, namely $\star|_{U^{p,q}} = i^{p+q}$. If we plot the (nontrivial) spaces $U^{p,q}$ in a lattice, each diagonal is made either of SD- or ASD-forms, with $U^{n,0}$ made of SD-forms (see Figure 1).




Figure 1: Representation of the spaces of SD and ASD in terms of the (p, q) -decomposition of forms on a 4-dimensional generalized Hermitian structure.

Definition 1.7. A *positive $U(n)$ structure* or a *positive Hermitian structure* on V is a generalized metric \mathcal{G} and a complex structure \mathcal{I}_+ on V_+ , the $+1$ -eigenspace of \mathcal{G} , orthogonal with respect to the natural pairing. A *negative $U(n)$ structure* or *negative Hermitian structure* on V is a generalized metric with an orthogonal complex structure \mathcal{I}_- on its -1 -eigenspace. We say that a generalized complex structure \mathcal{J} *extends* a positive/negative $U(n)$ structure $(\mathcal{G}, \mathcal{I})$ if \mathcal{I} is the restriction of \mathcal{J} to the appropriate space and $(\mathcal{G}, \mathcal{J})$ is a generalized Hermitian structure.

Given a generalized Hermitian structure, since \mathcal{J}_1 and \mathcal{G} commute, \mathcal{J}_1 preserves the eigenspaces of \mathcal{G} and hence, upon restriction to V_{\pm} , one obtains a positive and a negative Hermitian structure. Conversely, a positive (resp. negative) Hermitian structure can be extended to \mathbb{V} by declaring that it vanishes on V_- (resp. V_+). Then a pair of positive and negative Hermitian structures, \mathcal{I}_+ , \mathcal{I}_- gives rise to a generalized Hermitian structure by declaring that $\mathcal{J}_1 = \mathcal{I}_+ + \mathcal{I}_-$.

Given a positive $U(n)$ structure on V , we can use the isomorphism $V_+ \cong V$ to transport the metric and the complex structure \mathcal{I}_+ from V_+ to V , making it into a Hermitian vector space (V, g, I) . Further, we can use \mathcal{I}_+ to define a complex structure \mathcal{I}_- on V_- using the isomorphisms $V_+ \cong V \cong V_-$ and this way we have an extension of the $U(n)$ structure to a generalized Hermitian structure: namely we declare that \mathcal{J}_1 is \mathcal{I}_+ on V_+ and \mathcal{I}_- on V_- , hence, in the metric splitting of \mathbb{V} , \mathcal{J}_1 is the generalized complex structure associated to the complex structure² I and consequently \mathcal{J}_2 is the generalized complex structure associated to the Hermitian form $\omega = g \circ I$:

$$\mathcal{J}_1 = \begin{pmatrix} I & 0 \\ 0 & -I^* \end{pmatrix} \quad \mathcal{J}_2 = \begin{pmatrix} 0 & -\omega^{-1} \\ \omega & 0 \end{pmatrix}. \quad (1.6)$$

For this set of choices, there is a relation between the (p, q) -decomposition of forms determined by the generalized Hermitian structure and the usual (p, q) -decomposition of forms determined by the complex structure I on V .

Proposition 1.8. (Cavalcanti [7]) *Let \mathcal{J}_1 and \mathcal{J}_2 be as above and let L_2 be the $+i$ -eigenspace of \mathcal{J}_2 . The map*

$$\psi : \mathbb{V}_{\mathbb{C}} \longrightarrow \mathbb{V}_{\mathbb{C}}; \quad \psi(X + \xi) = e^{i\omega} e^{\frac{i}{2}\omega^{-1}} (X + \xi)$$

preserves \mathcal{J}_1 and

$$\psi(V_{\mathbb{C}}) = L_2, \quad \psi(V_{\mathbb{C}}^*) = \overline{L_2}.$$

Therefore the corresponding action on spinors,

$$\Psi : \wedge^{\bullet} V_{\mathbb{C}}^* \longrightarrow \wedge^{\bullet} V_{\mathbb{C}}^*; \quad \Psi(\varphi) = e^{i\omega} e^{\frac{i}{2}\omega^{-1}} (\varphi),$$

preserves the eigenspaces of \mathcal{J}_1 and maps $\wedge^k V_{\mathbb{C}}^$ into $U_{\mathcal{J}_2}^{n-k}$ that is*

$$\Psi(\wedge^{p,q} V^*) = U_{\mathcal{J}_1, \mathcal{J}_2}^{q-p, n-p-q}.$$

A positive Hermitian structure is fully determined by the $+i$ -eigenspace of \mathcal{I}_+ , that is, an isotropic n -dimensional subspace $V_+^{1,0} \subset \mathbb{T}_{\mathbb{C}}M$ for which $V_+^{1,0} \cap \overline{V_+^{1,0}} = \{0\}$ and $\langle v, \bar{v} \rangle > 0$ for all $v \in V_+^{1,0} \setminus \{0\}$.

If we are given a generalized complex extension \mathcal{J} of a positive Hermitian structure $(\mathcal{G}, \mathcal{I}_+)$ we obtain a bigrading of forms into $U^{p,q}$ as explained earlier. However, from the point of view of the $U(n)$ structure, the natural spaces to consider are $W_+^k = \sum_{p+q=k} U^{p,q}$. Indeed W_+^n corresponds to the vector space of all forms that are annihilated by the Clifford action of $V_+^{1,0}$ and $W_+^{n-2k} = \wedge^k V_+^{0,1} \cdot W_+^n$, that is, the spaces W_+^k are solely determined by the complex structure \mathcal{I}_+ . Notice that W_+^k is only nontrivial if $-n \leq k \leq n$ and $n - k = 0 \pmod{2}$. Further, since the spaces W_+^k are the diagonals of the $U^{p,q}$ decomposition, each W_+^k is made of either SD- or ASD-forms, with W_+^n being SD.

Another description of the spaces W_+^{\bullet} is obtained by extending \mathcal{I}_+ to an endomorphism of $\mathbb{T}M$ by declaring that $\mathcal{I}_+|_{V_-}$ vanishes, so that \mathcal{I}_+ is a skew-symmetric operator on $\mathbb{T}M$ with respect to the natural pairing, that is, $\mathcal{I}_+ \in \mathfrak{spin}(\mathbb{T}M)$. Similarly to a generalized complex structure, \mathcal{J} , for which the space U^k is the ik -eigenspace of the action of \mathcal{J} , letting \mathcal{I}_+ act on forms one sees that the space W_+^k is the $i\frac{k}{2}$ -eigenspace of \mathcal{I}_+ . Hence, Lemma 1.6 takes the following form for positive $U(n)$ structures:

²Note that usually, the generalized complex structure associated to a given a complex structure I on a vector space differs from \mathcal{J}_1 by a sign, or said another way, by overall conjugation. Hence, all the usual concepts get conjugated with this definition, for example, later we will have $\partial_{\mathcal{J}_1} = \bar{\partial}$ and the generalized canonical bundle is $\wedge^{0,n} T^*M$

Lemma 1.9. Let $(\mathcal{G}, \mathcal{I}_+)$ be a positive Hermitian structure on V , and let $\mathbb{I}_+ = e^{\frac{\pi \mathcal{I}_+}{2}}$. Then

$$\mathbb{I}_+^2 = -\star,$$

that is $\star|_{W_\pm^k} = -i^k$.

Similarly, for a negative Hermitian structure $(\mathcal{G}, \mathcal{I}_-)$, we can extend \mathcal{I}_- to \mathbb{V} by declaring that it vanishes on V_+ . If \mathcal{J}_1 is a generalized complex extension of \mathcal{I}_- , then the eigenspaces of \mathcal{I}_- correspond to the anti-diagonals of the $U^{p,q}$ decomposition: $W_-^k = \sum_{p-q=k} U^{p,q}$.

Finally we observe that the spaces W_\pm^k have both even and odd forms, so one can refine this grading to a $\mathbb{Z} \times \mathbb{Z}_2$ -grading:

$$W_\pm^{k,0} = W_\pm^k \cap (\wedge^{ev} V^* \otimes \mathbb{C}), \quad W_\pm^{k,1} = W_\pm^k \cap (\wedge^{od} V^* \otimes \mathbb{C}). \quad (1.7)$$

In what follows we will refer to both spaces W_\pm^k and $W_\pm^{k,l}$, with the understanding that if the \mathbb{Z}_2 -grading is not particularly important, we will simply omit it.

2 Intrinsic torsion of generalized Hermitian structures

Except for a generalized metric, each of the structures introduced in Section 1 has an appropriate integrability condition. We let (M^{2n}, H) be a manifold with a real closed 3-form H and consider the Courant bracket on sections of $\mathbb{T}M = TM \oplus T^*M$:

$$\llbracket X + \xi, Y + \eta \rrbracket_H = [X, Y] + \mathcal{L}_X \eta - i_Y d\xi - i_Y i_X H.$$

We will omit the 3-form from the bracket if it is clear from the context.

The Courant bracket is the derived bracket associated to the operator $d^H = d + H \wedge$, i.e., the following identity holds for all $v_1, v_2 \in \Gamma(\mathbb{T}M)$ and $\varphi \in \Omega^\bullet(M)$:

$$\begin{aligned} \llbracket v_1, v_2 \rrbracket_H \cdot \varphi &= \{\{v_1, d^H\}, v_2\} \varphi \\ &= d^H(v_1 \cdot v_2 \cdot \varphi) + v_1 \cdot d^H(v_2 \cdot \varphi) - v_2 \cdot d^H(v_1 \cdot \varphi) - v_2 \cdot v_1 \cdot d^H \varphi, \end{aligned} \quad (2.1)$$

where \cdot denotes the Clifford action of $\text{Clif}(\mathbb{T}M)$ on $\wedge^\bullet T^*M$ and $\{\cdot, \cdot\}$ denotes the graded commutator of operators.

The orthogonal action of a 2-form $B \in \Omega^2(M)$ on $\mathbb{T}M$ relates different Courant brackets:

$$\llbracket e_*^B v_1, e_*^B v_2 \rrbracket_H = e_*^B \llbracket v_1, v_2 \rrbracket_{H+dB}.$$

Definition 2.1. For each of the structures introduced in Definitions 1.5 and 1.7, we refer to the smooth assignment of such structure to $T_x M$ for each $x \in M$ by including the adjective *almost* in the name of the structure.

Definition 2.2 (Integrability conditions).

- A (*integrable*) *generalized complex structure* is a generalized almost complex structure \mathcal{J} whose $+i$ -eigenspace is involutive with respect to the Courant bracket.
- A *generalized Hermitian structure* is a pair $(\mathcal{G}, \mathcal{J}_1)$ of generalized metric and compatible integrable generalized complex structure.
- A *generalized Kähler structure* is a generalized Hermitian structure $(\mathcal{G}, \mathcal{J}_1)$ for which \mathcal{J}_1 and $\mathcal{J}_2 = \mathcal{J}_1 \mathcal{G}$ are integrable.

2.1 The Nijenhuis tensor

Let us spend some time to understand the *Nijenhuis tensor* of an almost generalized complex structure \mathcal{J} . This tensor is defined in the usual way, namely if L is the $+i$ -eigenspace of \mathcal{J}

$$\text{Nij} : \Gamma(\bar{L}) \times \Gamma(\bar{L}) \longrightarrow \Gamma(L); \quad \text{Nij}(X, Y) = -\llbracket X, Y \rrbracket^L, \quad (2.2)$$

where \cdot^L indicates projection onto L . We can alternatively use the identification $L = \bar{L}^*$ from (1.4) to consider the operator

$$\begin{aligned} N : \Gamma(\bar{L}) \times \Gamma(\bar{L}) \times \Gamma(\bar{L}) &\longrightarrow \Omega^0(M; \mathbb{C}); \\ N(X, Y, Z) &= -2\langle \llbracket X, Y \rrbracket, Z \rangle = 2\langle \text{Nij}(X, Y), Z \rangle. \end{aligned} \quad (2.3)$$

As usual, Nij is a tensor, indeed, for $f \in \mathbb{C}^\infty(M; \mathbb{C})$ we have

$$\text{Nij}(X, fY) = -\llbracket X, fY \rrbracket^L = -(f\llbracket X, Y \rrbracket + (\mathcal{L}_{\pi_T(X)}f)Y)^L = -f\llbracket X, Y \rrbracket^L = f\text{Nij}(X, Y).$$

Further, the tensor $N \in \wedge^2 L \otimes L$ defined above actually lies in $\wedge^3 L$. Indeed, for $X, Y, Z \in \Gamma(\bar{L})$ we have

$$0 = \mathcal{L}_{\pi_T(X)}\langle Y, Z \rangle = -(\langle \llbracket X, Y \rrbracket, Z \rangle + \langle Y, \llbracket X, Z \rrbracket \rangle) = \frac{1}{2}(N(X, Y, Z) + N(X, Z, Y)),$$

which shows that N is fully skew.

A different way to understand N arises by using the U^k decomposition of forms determined by \mathcal{J} . Namely, letting $\mathcal{U}^k = \Gamma(U^k)$ (throughout the paper we denote the sheaf of sections of U^k by \mathcal{U}^k and the sheaf of sections of $U^{p,q}$ by $\mathcal{U}^{p,q}$), one has:

Lemma 2.3. In an almost generalized complex manifold

$$d^H : \mathcal{U}^k \longrightarrow \mathcal{U}^{k-3} \oplus \mathcal{U}^{k-1} \oplus \mathcal{U}^{k+1} \oplus \mathcal{U}^{k+3}. \quad (2.4)$$

Further the map $\pi_{k+3} \circ d^H : \mathcal{U}^k \longrightarrow \mathcal{U}^{k+3}$ corresponds to the Clifford action of N , the Nijenhuis tensor defined in (2.3).

Proof. We prove first that

$$d^H : \mathcal{U}^k \longrightarrow \sum_{j \geq k-3} \mathcal{U}^j \quad (2.5)$$

and will do so by induction, starting at $k = n + 1$ and working our way down. For $k = n + 1$ we have that $U^{n+1} = \{0\}$ and the claim follows trivially.

Next we assume the result to be true for all $j > k$ and let $\rho \in \mathcal{U}^k$. For $v_1, v_2 \in \Gamma(L)$, using (2.1) we have

$$v_2 \cdot v_1 \cdot d^H \rho = -\llbracket v_1, v_2 \rrbracket \cdot \rho + d^H(v_1 \cdot v_2 \cdot \rho) + v_1 d^H(v_2 \cdot \rho) - v_2 d^H(v_1 \cdot \rho).$$

Since the Clifford action of v_i sends \mathcal{U}^j to \mathcal{U}^{j+1} , the inductive hypothesis implies that the last three terms lie in $\oplus_{j \geq k-1} \mathcal{U}^j$, while the first term, being the action of an element of $L \oplus \bar{L}$ on ρ lies in $\mathcal{U}^{k+1} \oplus \mathcal{U}^{k-1}$. Therefore we conclude that

$$v_2 \cdot v_1 \cdot d^H \rho \in \oplus_{j \geq k-1} \mathcal{U}^j \quad \text{for all } v_1, v_2 \in \Gamma(L)$$

and (2.5) follows.

As d^H is a real operator and $U^{-k} = \overline{U^k}$ conjugating (2.5) we have that

$$d : \mathcal{U}^k \longrightarrow \sum_{j \leq k-3} \mathcal{U}^j.$$

Furthermore, if U^k is made of even forms then U^{k+1} is made of odd forms and vice versa, we have that

$$d^H : \mathcal{U}^k \longrightarrow \sum_{\substack{|j-k| \leq 3 \\ j-k=1 \pmod{2}}} \mathcal{U}^j,$$

therefore proving (2.4).

Now we prove that $\pi_{k+3} \circ d^H$ corresponds to the Clifford action of N . Once again we use induction, this time starting at $k = -n - 1$ and moving upwards. Since $\mathcal{U}^{-n-1} = \{0\}$, the claim is trivial there. Assume now that for $j < k$ we have proved that $\pi_{j+3} \circ d^H$ is the Clifford action of N . Let $\rho \in \mathcal{U}^k$ and $v_1, v_2 \in \Gamma(\bar{L})$. Then using (2.1) we have

$$\begin{aligned} v_2 v_1 \pi_{k+3} \circ d^H \rho &= \pi_{k+1}(v_2 v_1 d^H \rho) \\ &= \pi_{k+1}(-\llbracket v_1, v_2 \rrbracket \rho + d^H v_1 v_2 \rho + v_1 d^H v_2 \rho - v_2 d^H v_1 \rho) \\ &= N(v_1, v_2) \rho + \pi_{k+1} \circ d^H(v_1 v_2 \rho) + v_1 \pi_{k+2} \circ d^H v_2 \rho - v_2 \pi_{k+2} \circ d^H v_1 \rho \\ &= \{v_2, \{v_1, N\}\} \rho + N v_1 v_2 \rho + v_1 N v_2 \rho - v_2 N v_1 \rho \\ &= v_2 v_1 N \rho, \end{aligned}$$

where in third equality we have used that the component of $\llbracket v_1, v_2 \rrbracket \rho$ in \mathcal{U}^{k+1} is given by the Clifford action of the L component of $\llbracket v_1, v_2 \rrbracket$, that is $\iota_{v_2} \iota_{v_1} N$ and in the fourth equality we used the inductive hypothesis as well as the fact that when acting on forms, the interior product of $v \in \Gamma(\bar{L})$ with $\varphi \in \Gamma(\wedge^\bullet L)$ is given by $\varphi(v) \rho = \{v, \varphi\} \rho$. The last equality follows by expanding the graded commutator and canceling out similar terms. \square

If we compose $d^H|_{\mathcal{U}^k}$ with projection onto \mathcal{U}^{k+1} and \mathcal{U}^{k-1} we get operators

$$\partial_{\mathcal{J}} = \pi_{k+1} \circ d^H : \mathcal{U}^k \longrightarrow \mathcal{U}^{k+1} \quad \text{and} \quad \bar{\partial}_{\mathcal{J}} = \pi_{k-1} \circ d^H : \mathcal{U}^k \longrightarrow \mathcal{U}^{k-1},$$

and if the generalized complex structure \mathcal{J} is clear from the context, we denote these operators simply by ∂ and $\bar{\partial}$. As proved by Gualtieri in [21], integrability is equivalent to the requirement

$$d^H : \mathcal{U}^k \longrightarrow \mathcal{U}^{k-1} \oplus \mathcal{U}^{k+1}; \quad d^H = \partial + \bar{\partial},$$

so we can also see from this point of view how the vanishing of the Nijenhuis tensor implies integrability.

2.2 The intrinsic torsion and the road to integrability

With this understanding of the Nijenhuis tensor, we can give a pictorial description of the long road to integrability from almost generalized Hermitian to generalized Kähler. Indeed, given an almost generalized Hermitian structure, we get a splitting of forms into spaces $\mathcal{U}^{p,q}$. According to Lemma 2.3, d^H can not change either the ‘ p ’ or the ‘ q ’ grading by more than three and it must switch parity. Hence d^H decomposes as a sum of eight operators and their complex conjugates

$$\begin{aligned} d^H &= \delta_+ + \bar{\delta}_+ + \delta_- + \bar{\delta}_- + N_+ + \bar{N}_+ + N_- + \bar{N}_- + N_1 + \bar{N}_1 + N_2 + \bar{N}_2 + N_3 + \bar{N}_3 + N_4 + \bar{N}_4; \\ \delta_+ : \mathcal{U}^{p,q} &\longrightarrow \mathcal{U}^{p+1,q+1}, & \delta_- : \mathcal{U}^{p,q} &\longrightarrow \mathcal{U}^{p+1,q-1}, \\ N_+ : \mathcal{U}^{p,q} &\longrightarrow \mathcal{U}^{p+3,q+3}, & N_- : \mathcal{U}^{p,q} &\longrightarrow \mathcal{U}^{p+3,q-3}, \\ N_1 : \mathcal{U}^{p,q} &\longrightarrow \mathcal{U}^{p-1,q+3}, & N_2 : \mathcal{U}^{p,q} &\longrightarrow \mathcal{U}^{p+1,q+3}, \\ N_3 : \mathcal{U}^{p,q} &\longrightarrow \mathcal{U}^{p+3,q+1}, & N_4 : \mathcal{U}^{p,q} &\longrightarrow \mathcal{U}^{p+3,q-1}. \end{aligned} \tag{2.6}$$

and we can draw in a diagram all the possible nontrivial $\mathcal{U}^{p,q}$ components of $d^H|_{\mathcal{U}^{p,q}}$ as arrows (see Figure 2).




Figure 2: Representation of the nontrivial components of d^H when restricted to $\mathcal{U}^{p,q}$ for a generalized almost Hermitian structure.

Once we require that \mathcal{J}_1 is integrable, i.e. we are dealing in fact with a generalized Hermitian structure, then d^H only changes the ‘ p ’ degree by ± 1 and several components of d^H present in the nonintegrable case, now vanish and the diagram from Figure 2 clears up to the one presented in Figure 3.




Figure 3: Representation of the nontrivial components of d^H for a generalized Hermitian structure.

Finally, if we require that \mathcal{J}_2 is also integrable, and hence we are in fact dealing with a generalized Kähler structure, the last two components of the Nijenhuis tensor, labeled N_1 and N_2 above, vanish and d^H decomposes as a sum of four operators, as pictured in Figure 4.

This shows that the obstruction for a generalized almost Hermitian structure to be a generalized Kähler structure is given by the tensors N_i , $i = 1, 2, 3, 4$ and N_{\pm} .

Definition 2.4. The *intrinsic torsion* of a generalized Hermitian manifold are the tensors N_i , $i = 1, 2, 3, 4$ and N_{\pm} .

Next sections we will introduce geometric structures on M which are weaker than generalized Kähler structures and show how these structures can be phrased in terms of the vanishing of certain components of the intrinsic torsion.




Figure 4: Representation of the nontrivial components of d^H for a generalized Kähler structure.

2.3 The operators δ_{\pm} and $\bar{\delta}_{\pm}$

Not requiring integrability, d^H restricted to $\mathcal{U}^{p,q}$ has sixteen components but only four are not tensorial, namely δ_{\pm} and $\bar{\delta}_{\pm}$. Hence we have

$$d^H \sim \delta_+ + \bar{\delta}_+ + \delta_- + \bar{\delta}_-, \quad (2.7)$$

where \sim indicates that these operators agree up to lower order terms, i.e., they have the same symbol. Using the decomposition

$$\mathbb{T}_{\mathbb{C}}M = V_+^{1,0} \oplus V_+^{0,1} \oplus V_-^{1,0} \oplus V_-^{0,1}, \quad (2.8)$$

one can easily see that the decomposition (2.7) in terms of symbols corresponds to the decomposition of a 1-form $\xi \in \mathbb{T}M$ into its four components according to (2.8) and one can also check that the symbol sequence for each of the operators δ_{\pm} and $\bar{\delta}_{\pm}$ is exact, e.g., for δ_+ , the symbol sequence associated to sequence of operators

$$\dots \xrightarrow{\delta_+} \mathcal{U}^{p-1,q-1} \xrightarrow{\delta_+} \mathcal{U}^{p,q} \xrightarrow{\delta_+} \mathcal{U}^{p+1,q+1} \xrightarrow{\delta_+} \dots$$

is an exact sequence. Adding over $p+q=k$ and letting $\mathcal{W}_+^k = \bigoplus_{p+q=k} \mathcal{U}^{p,q}$ we have also that the symbol sequence associated to

$$\dots \xrightarrow{\delta_+} \mathcal{W}_+^{k-2} \xrightarrow{\delta_+} \mathcal{W}_+^k \xrightarrow{\delta_+} \mathcal{W}_+^{k+2} \xrightarrow{\delta_+} \dots \quad (2.9)$$

is exact. And similarly, adding over q we get that the symbol sequence of

$$\dots \xrightarrow{\delta_+} \mathcal{U}^{p-1} \xrightarrow{\delta_+} \mathcal{U}^p \xrightarrow{\delta_+} \mathcal{U}^{p+1} \xrightarrow{\delta_+} \dots \quad (2.10)$$

is exact.

3 Parallel Hermitian and bi-Hermitian structures

The first type of structures that we will relate to the intrinsic torsion are Hermitian structures which are parallel for a connection with closed skew torsion. The existence of a relationship between connections with closed, skew symmetric torsion and the Courant bracket was made evident by Hitchin in [23]. Precisely, given a generalized metric on $\mathbb{T}M$, we let H be the 3-form corresponding to the metric splitting and g be the induced metric on M . Also, for $X \in \Gamma(TM)$ we let $X_{\pm} \in \Gamma(V_{\pm})$ be unique lifts of X to V_{\pm} , we let $\pi_{\pm} : \mathbb{T}M \rightarrow V_{\pm}$ be the orthogonal projections onto V_{\pm} and $\pi_T : \mathbb{T}M \rightarrow TM$ be the natural projection.

Proposition 3.1 (Hitchin [23]). *Let $\tilde{\nabla}^\pm$ be the unique metric connection whose torsion is skew symmetric and equal to $\mp H$. Then*

$$\tilde{\nabla}_X^+ Y = \pi_T \circ \pi_+ \llbracket X_-, Y_+ \rrbracket,$$

$$\tilde{\nabla}_X^- Y = \pi_T \circ \pi_- \llbracket X_+, Y_- \rrbracket.$$

From this proposition, we see that the isomorphisms $\pi_T : V_\pm \rightarrow TM$ relate the connections with torsion $\mp H$ to the operators

$$\nabla^+ : \Gamma(V_-) \times \Gamma(V_+) \rightarrow \Gamma(V_+); \quad \nabla_v^+ w = \pi_+ \llbracket v, w \rrbracket, \quad v \in \Gamma(V_+), w \in \Gamma(V_-), \quad (3.1)$$

$$\nabla^- : \Gamma(V_+) \times \Gamma(V_-) \rightarrow \Gamma(V_-); \quad \nabla_w^- v = \pi_- \llbracket w, v \rrbracket, \quad v \in \Gamma(V_+), w \in \Gamma(V_-). \quad (3.2)$$

As we will work with the spaces V_\pm directly, we will use ∇^\pm instead of the connections they induce on TM , with the understanding that these are equivalent operators, so, for example, if $\tilde{\nabla}^+$ has holonomy in $U(n)$, then M has an almost Hermitian structure (g, I) which is parallel with respect to $\tilde{\nabla}^+$ and via the isomorphism $\pi_T : V_+ \rightarrow TM$, V_+ gets a positive almost Hermitian structure which is parallel for ∇^+ so that ∇^+ has holonomy in $U(n)$ and the converse statement also holds.

Definition 3.2. *A parallel positive (resp. negative) Hermitian structure is a positive (resp. negative) almost Hermitian structure which is parallel with respect to ∇^+ , (resp. ∇^-). A parallel bi-Hermitian structure or parallel $U(n) \times U(n)$ -structure is a triple (g, I_+, I_-) such that (g, I_+) is a parallel positive Hermitian structure and (g, I_-) is a parallel negative Hermitian structure.*

Proposition 3.3. *If the connection ∇^+ has holonomy in $U(n)$, the corresponding almost Hermitian structure satisfies*

$$\llbracket \Gamma(V_+^{1,0}), \Gamma(V_+^{1,0}) \rrbracket \subset \Gamma(V_+ \otimes \mathbb{C}). \quad (3.3)$$

And conversely, a Hermitian structure satisfying (3.3) is parallel with respect to ∇^+ .

Proof. Let \mathcal{I}_+ be a complex structure on V_+ orthogonal with respect to the natural pairing and let $V_+^{1,0}$ be its $+i$ -eigenspace. Then, for $v_1, v_2 \in \Gamma(V_+^{1,0})$ and $w \in \Gamma(V_-)$, we have that $\langle v_i, w \rangle = 0$, as V_+ and V_- are orthogonal with respect to the natural pairing. Hence

$$\begin{aligned} 0 &= \mathcal{L}_{\pi_T v_1} \langle w, v_2 \rangle \\ &= \langle \llbracket v_1, w \rrbracket, v_2 \rangle + \langle w, \llbracket v_1, v_2 \rrbracket \rangle \\ &= -\langle \llbracket w, v_1 \rrbracket, v_2 \rangle + \langle w, \llbracket v_1, v_2 \rrbracket \rangle \\ &= -\langle \nabla_w^+ v_1, v_2 \rangle + \langle w, \llbracket v_1, v_2 \rrbracket \rangle. \end{aligned} \quad (3.4)$$

where in the third equality we used again that V_+ and V_- are orthogonal with respect to the natural pairing, and hence $\llbracket v_1, w \rrbracket = -\llbracket w, v_1 \rrbracket$ and in the fourth equality we used that $v_2 \in V_+$ and hence the V_- component of $\llbracket w, v_1 \rrbracket$ is annihilated by the natural pairing.

If \mathcal{I}_+ is parallel with respect to ∇^+ , then $\langle \nabla_w^+ v_1, v_2 \rangle$ vanishes for all $v_1, v_2 \in \Gamma(V_+^{1,0})$ and $w \in \Gamma(V_-)$ and hence, according to (3.4), so does $\langle w, \llbracket v_1, v_2 \rrbracket \rangle$, showing that $\llbracket v_1, v_2 \rrbracket$ must be orthogonal to V_- and hence, a section of V_+ . Conversely, if $\llbracket v_1, v_2 \rrbracket \in \Gamma(V_+)$, then for all $w \in \Gamma(V_-)$, $\langle w, \llbracket v_1, v_2 \rrbracket \rangle = 0$ hence (3.4) implies that $\langle \nabla_w^+ v_1, v_2 \rangle = 0$ showing that $\nabla_w^+ v_1$ is orthogonal to $V_+^{1,0}$. Since $V_+^{1,0}$ is a maximal isotropic of $V_+ \otimes \mathbb{C}$, we conclude that $\nabla_w^+ v_1 \in \Gamma(V_+^{1,0})$ for all $w \in V_-$ and $v_1 \in \Gamma(V_+^{1,0})$, hence \mathcal{I}_+ is parallel with respect to ∇^+ . \square

Obviously the same result holds exchanging ∇^+ and $V_+^{1,0}$ by ∇^- and $V_-^{1,0}$.

Next, and throughout these notes, we denote by \mathcal{W}^k and $\mathcal{W}^{k,l}$ the sheaf of sections of the bundles W^k and $W^{k,l}$ respectively.

Theorem 3.4. Let (M, H) be a manifold with 3-form, \mathcal{G} be a generalized metric on M and $\mathcal{I}_+, \mathcal{I}_-$ be a positive and a negative almost Hermitian structure on M

- \mathcal{I}_+ is parallel with respect to ∇^+ if and only if $N_2 = N_3 = 0$, i.e.,

$$d^H : \mathcal{W}_+^k \longrightarrow \mathcal{W}_+^{k-6} \oplus \mathcal{W}_+^{k-2} \oplus \mathcal{W}_+^k \oplus \mathcal{W}_+^{k+2} \oplus \mathcal{W}_+^{k+6};$$

- \mathcal{I}_- is parallel with respect to ∇^- if and only if $N_1 = N_4 = 0$, i.e.,

$$d^H : \mathcal{W}_-^k \longrightarrow \mathcal{W}_-^{k-6} \oplus \mathcal{W}_-^{k-2} \oplus \mathcal{W}_-^k \oplus \mathcal{W}_-^{k+2} \oplus \mathcal{W}_-^{k+6};$$

- $(\mathcal{G}, \mathcal{I}_+, \mathcal{I}_-)$ is a parallel bi-Hermitian structure if and only if $N_i = 0$ for $i = 1, 2, 3, 4$.

Proof. Since V_- is the orthogonal complement of V_+ with respect to the natural pairing, it is clear that (3.3) is equivalent to the following two conditions

$$\langle \llbracket v_1, v_2 \rrbracket, w \rangle = 0 \text{ for all } v_1, v_2 \in \Gamma(V_+^{1,0}), w \in \Gamma(V_-^{0,1}).$$

$$\langle \llbracket v_1, v_2 \rrbracket, w \rangle = 0 \text{ for all } v_1, v_2 \in \Gamma(V_+^{1,0}), w \in \Gamma(V_-^{1,0});$$

but the first condition is equivalent to the vanishing of \overline{N}_2 and the second, to the vanishing of \overline{N}_3 . Finally, since the component of d^H mapping \mathcal{W}^k to \mathcal{W}^{k+4} is given by the sum $N_2 + N_3$, we see that the vanishing of this component is also equivalent to the parallel condition.

The remaining claims are proved similarly. \square

The decomposition of d^H for an almost generalized Hermitian structure extending a parallel positive Hermitian structure is depicted in Figure 5 and the decomposition of d^H for a parallel bi-Hermitian structure is depicted in Figure 6.




Figure 5: Representation of the nontrivial components of d^H for an almost generalized Hermitian structure extending a parallel positive Hermitian structure

4 SKT structures

Classically, an SKT structure is a Hermitian structure (M, g, I) for which $dd^c\omega = 0$, where ω is the Hermitian 2-form. In this case, $d^c\omega = H$ is a closed 3-form and the complex structure is in fact parallel with respect to the metric connection with torsion $-H$, hence SKT structures are a




Figure 6: Representation of the nontrivial components of d^H for a parallel bi-Hermitian structure.

particular case of positive parallel Hermitian structures, the only difference being that now one requires I to be integrable. In this section we phrase the SKT condition in terms of generalized geometry and relate it with the intrinsic torsion. We will see that an SKT structure lies precisely half way between a generalized Hermitian and a generalized Kähler structure.

Given an SKT structure (g, I) we let H be the background 3-form and consider the generalized metric \mathcal{G} as in (1.2). Using the isomorphism $V_+ \cong TM$ we use I to induce a complex structure \mathcal{I}_+ on V_+ and hence split V_+ into eigenspaces of \mathcal{I}_+ : $V_+ \otimes \mathbb{C} = V_+^{1,0} \oplus V_+^{0,1}$.

Proposition 4.1. (Cavalcanti [8]) *For an SKT structure (g, I) on M with $d^c\omega = H$, $V_+^{1,0}$ is involutive with respect to the H -Courant bracket.*

Conversely, given a positive almost Hermitian structure $(\mathcal{G}, \mathcal{I}_+)$ on (M, H) , where H is the 3-form associated of the metric splitting of TM , if $V_+^{1,0}$ is involutive, then the induced Hermitian structure (g, I) on M is an SKT structure with $d^c\omega = H$.

Remark. Here we obtain, in a new light, a well known contrast between connections with torsion and the Levi-Civita connection regarding integrability. Indeed, for the Levi-Civita connection, reduction of the holonomy group to $U(n)$ implies integrability of the complex structure, but that is known not to be the case for connections with torsion. From our point of view, this is the difference between the reduced holonomy condition

$$[\Gamma(V_+^{1,0}), \Gamma(V_+^{1,0})] \subset \Gamma(V_+)$$

and the SKT condition

$$[\Gamma(V_+^{1,0}), \Gamma(V_+^{1,0})] \subset \Gamma(V_+^{1,0}).$$

Of course, if we have $d^c\omega = -H$, we can lift the SKT data to V_- to obtain a negative Hermitian structure and an analogous version of Proposition 4.1 holds.

This result motivates our formulation of the SKT condition.

Definition 4.2. A *positive (resp. negative) SKT structure* is a positive (resp. negative) almost $U(n)$ structure $(\mathcal{G}, \mathcal{I})$ for which the $+i$ -eigenspace of \mathcal{I} is involutive. A generalized (almost) complex structure \mathcal{J} *extends* an SKT structure if \mathcal{J} is fiberwise an extension of \mathcal{I} , in which case we say that \mathcal{J} is a *generalized (almost) complex/Hermitian extension* of the SKT structure.

Firstly, we observe that the SKT condition can also be phrased in terms of the vanishing of components of the intrinsic torsion. Indeed, in the presence of an extension of, say, a positive $U(n)$

structure $(\mathcal{G}, \mathcal{I}_+)$ to a generalized almost complex structure \mathcal{J}_1 if we let $V_+^{1,0}$ be the $+i$ -eigenspace of \mathcal{I}_+ then involutivity is equivalent to

$$\langle [v_1, v_2], w_1 \rangle = 0 \quad \text{for all } v_1, v_2 \in \Gamma(V_+^{1,0}), w \in \Gamma(V_+^{1,0} \oplus V_-),$$

that is, the components N_2, N_3 and N_+ of the intrinsic torsion vanish. A similar argument gives a characterization of negative SKT structures.

Proposition 4.3. *Let \mathcal{G} be a generalized metric on a manifold with 3-form (M, H) and let \mathcal{I}_\pm be a pair of positive and negative Hermitian structures compatible with \mathcal{G} . If $\mathcal{J}_1 = \mathcal{I}_+ + \mathcal{I}_-$ then $(\mathcal{G}, \mathcal{I}_+)$ is an SKT structure if and only if N_2, N_3 and N_+ vanish and $(\mathcal{G}, \mathcal{I}_-)$ is a negative SKT structure if and only if N_1, N_4 and N_- vanish.*




Figure 7: Representation of the nontrivial components of d^H for a positive SKT structure

Corollary 4.4. Let (M, H) be a manifold with 3-form.

- A parallel positive (resp. negative) almost Hermitian structure is a positive (resp. negative) SKT structure if and only if $N_+ = 0$ (resp. $N_- = 0$);
- A parallel bi-Hermitian structure is a generalized Kähler structure if and only if $N_+ = N_- = 0$.

Corollary 4.5. Let M be a four dimensional manifold.

- A parallel positive/negative Hermitian structure is a positive/negative SKT structure;
- A parallel bi-Hermitian structure is a generalized Kähler structure.

Proof. Since M is four dimensional, $V_\pm^{1,0}$ are two dimensional complex vector spaces, so $\wedge^3 V_\pm^{1,0} = \{0\}$ and hence $N_\pm \in \wedge^3 V_\pm^{1,0}$ are the trivial tensors. \square

Corollary 4.6 (Gualtieri [20]). Let \mathcal{G} be a generalized metric on a manifold with 3-form (M, H) , \mathcal{I}_\pm be a pair of positive and negative SKT structures compatible with \mathcal{G} and $\mathcal{J}_1 = \mathcal{I}_+ + \mathcal{I}_-$. Then $(\mathcal{G}, \mathcal{J}_1)$ is a generalized Kähler structure.

Proof. According to Proposition 4.3, under the hypothesis, all components of the intrinsic torsion vanish hence $(\mathcal{G}, \mathcal{J}_1)$ is a generalized Kähler structure. \square

Next, we describe the integrability condition for an SKT structure in terms of the decomposition of forms into W_{\pm}^k and $U^{p,q}$ (for a fixed generalized complex extension \mathcal{J}_1) described in the previous section.

Theorem 4.7. *Let $(\mathcal{G}, \mathcal{I}_+)$ be a positive almost Hermitian structure on a manifold with 3-form (M^{2n}, H) . Then the following are equivalent:*

1. $(\mathcal{G}, \mathcal{I}_+)$ is a positive SKT structure;
2. $d^H : \mathcal{W}_+^k \longrightarrow \mathcal{W}_+^{k-2} \oplus \mathcal{W}_+^k \oplus \mathcal{W}_+^{k+2}$ for all k ;
3. $d^H : \mathcal{W}_+^n \longrightarrow \mathcal{W}_+^{n-2} \oplus \mathcal{W}_+^n$.

Proof. We will first prove that 1) implies 2). Let \mathcal{I}_- be any complex structure on V_- orthogonal with respect to the natural pairing so that \mathcal{G} and $\mathcal{J}_1 = \mathcal{I}_+ + \mathcal{I}_-$ form a generalized almost Hermitian structure. Then according to Proposition 4.3, N_2, N_3 and N_+ vanish. Then, we see that d^H splits into three components:

$$d^H = \delta_+^N + \overline{\delta_+^N} + \not\delta_- \quad (4.1)$$

$$\begin{aligned} \not\delta_- : \mathcal{W}_+^k &\longrightarrow \mathcal{W}_+^k; & \not\delta_- &= \delta_- + \overline{\delta_-} + N_- + \overline{N_-}, \\ \delta_+^N : \mathcal{W}_+^k &\longrightarrow \mathcal{W}_+^{k+2}; & \delta_+^N &= \delta_+ + N_1 + N_4, \\ \overline{\delta_+^N} : \mathcal{W}_+^k &\longrightarrow \mathcal{W}_+^{k-2}; & \overline{\delta_+^N} &= \overline{\delta_+} + \overline{N_1} + \overline{N_4}. \end{aligned}$$

Condition 2) clearly implies condition 3). Finally to prove that 3) implies 1) we once again choose a complex structure \mathcal{I}_- on V_- and observe that since $\overline{N_2}, \overline{N_3}$ and $\overline{N_+}$ are tensors, it is enough to check that they vanish when applied to spaces where their action is effective. But for $\varphi \in \mathcal{U}^{n,0} \subset \mathcal{W}_+^n$, the components of d^H landing on \mathcal{W}_+^{n-6} and \mathcal{W}_+^{n-4} are precisely $\overline{N_+} \cdot \varphi$ and $\overline{N_3} \cdot \varphi$ and the vanishing of these forms for $\varphi \neq 0$ implies that $\overline{N_+} = \overline{N_3} = 0$. Similarly, if $\psi \in \mathcal{U}^{0,n}$, the \mathcal{W}_+^{n-4} component of d^H is $\overline{N_2} \cdot \psi$, hence the vanishing of this component implies that $\overline{N_2} = 0$ and Proposition 4.3 implies that \mathcal{I}_+ is integrable. \square

While the different $U^{p,q}$ components of d^H obtained in terms of the generalized almost Hermitian extension of the SKT structure depend on the particular extension chosen, the operators $\delta_+^N, \overline{\delta_+^N}$ and $\not\delta_-$ depend only on the SKT data, since they are the decomposition of d^H obtained from the eigenspaces of \mathcal{I}_+ . Since $(d^H)^2 = 0$ these operators satisfy some relations.

Corollary 4.8. The following hold

$$(\delta_+^N)^2 = (\overline{\delta_+^N})^2 = 0; \quad \{\delta_+^N, \not\delta_-\} = 0; \quad \{\overline{\delta_+^N}, \not\delta_-\} = 0; \quad \{\delta_+^N, \overline{\delta_+^N}\} + (\not\delta_-^H)^2 = 0.$$

In the arguments up to now, given a positive SKT structure, we have chosen \mathcal{I}_- rather freely, but the complex structure \mathcal{I} corresponding to the SKT data (see Proposition 4.1) is integrable and together with \mathcal{G} forms a generalized Hermitian extension of $(\mathcal{G}, \mathcal{I}_+)$, that is, an SKT always has a generalized Hermitian extension.

Theorem 4.9. *Let (M, H) be a manifold with 3-form and $(\mathcal{G}, \mathcal{J}_1)$ be an generalized almost Hermitian structure on M . Then the following are equivalent:*

1. $(\mathcal{G}, \mathcal{J}_1)$ is an integrable generalized complex extension of a positive SKT structure;
2. $d^H : \mathcal{U}^{p,q} \longrightarrow \mathcal{U}^{p+1,q+1} \oplus \mathcal{U}^{p+1,q-1} \oplus \mathcal{U}^{p+1,q-3} \oplus \mathcal{U}^{p-1,q+3} \oplus \mathcal{U}^{p-1,q+1} \oplus \mathcal{U}^{p-1,q-1}$ for all $p, q \in \mathbb{Z}$;
3. $d^H : \mathcal{U}^{p,n-p} \longrightarrow \mathcal{U}^{p+1,n-p-1} \oplus \mathcal{U}^{p+1,n-p-3} \oplus \mathcal{U}^{p-1,n-p+1} \oplus \mathcal{U}^{p-1,n-p-1}$ for all $p \in \mathbb{Z}$;
4. $d^H : \mathcal{U}^{0,n} \longrightarrow \mathcal{U}^{-1,n-1} \oplus \mathcal{U}^{1,n-1} \oplus \mathcal{U}^{1,n-3}$ and $d^H : \mathcal{U}^{n,0} \longrightarrow \mathcal{U}^{n-1,1} \oplus \mathcal{U}^{n-1,-1}$.

Proof. The fact that 1) implies 2) follows from Proposition 4.3 and integrability of \mathcal{J}_1 . The implications 2) \Rightarrow 3) \Rightarrow 4) are immediate. Finally, similarly to the proof of Theorem 4.7, the values of the different components of the intrinsic torsion are fully determined by their action on $\mathcal{U}^{n,0}$ and $\mathcal{U}^{0,n}$ and 4) implies the vanishing of all components of the intrinsic torsion except from N_1 . \square

This theorem allows us to define six differential operators on a generalized complex extension of a positive SKT structure:

$$d^H = \delta_- + \bar{\delta}_- + N + \bar{N} + \delta_+ + \bar{\delta}_+ \quad (4.2)$$

$$\delta_- : \mathcal{U}^{p,q} \longrightarrow \mathcal{U}^{p+1,q-1}; \quad \delta_+ : \mathcal{U}^{p,q} \longrightarrow \mathcal{U}^{p+1,q+1}; \quad N : \mathcal{U}^{p,q} \longrightarrow \mathcal{U}^{p-1,q+3}.$$




Figure 8: Decomposition of d^H for a generalized complex extension of a positive SKT structure.

We see that $N = N_1$ is the Nijenhuis tensor of \mathcal{J}_2 and this theorem shows that if $(\mathcal{G}, \mathcal{J}_1)$ is a generalized Hermitian extension of an SKT structure then half of the Nijenhuis tensor of \mathcal{J}_2 vanishes, namely the component N_2 from (2.6). Also, this decomposition allows us to express the operators δ_+^N and δ_- from (4.1) in terms of δ_+ , δ_- , N and their conjugates:

$$\delta_+^N = \delta_+ + N; \quad \delta_- = \delta_- + \bar{\delta}_-. \quad (4.3)$$

Since $(d^H)^2 = 0$ these operators satisfy a number of relations.

Corollary 4.10. The following relations and their complex conjugates hold:

$$\delta_+^2 = N^2 = 0; \quad \{\delta_+, N\} = 0;$$

$$\delta_-^2 = -\{\bar{N}, \delta_+\}; \quad \{\bar{N}, \delta_-\} = 0; \quad \{\delta_-, \delta_+\} = 0; \quad \{\delta_+, \bar{\delta}_-\} = -\{\delta_-, N\};$$

$$\{\delta_+, \bar{\delta}_+\} + \{\delta_-, \bar{\delta}_-\} + \{N, \bar{N}\} = 0.$$

Corollary 4.11. Let $(\mathcal{G}, \mathcal{J}_1)$ be a generalized Hermitian structure on a manifold with 3-form (M, H) and let $\mathcal{J}_2 = \mathcal{G}\mathcal{J}_1$ be the associated generalized almost complex structure. If the canonical bundle of \mathcal{J}_2 admits a $\bar{\partial}_{\mathcal{J}_1}$ -closed trivialization, then $(\mathcal{G}, \mathcal{J})$ is an extension of an SKT structure.

Proof. Since \mathcal{J}_1 is integrable, $d^H : \mathcal{U}^{n,0} \longrightarrow \mathcal{U}^{n-1,1} \oplus \mathcal{U}^{n-1,-1}$. Due to Lemma 2.3, we have that

$$d^H : \mathcal{U}^{0,n} \longrightarrow \mathcal{U}^{-1,n-3} \oplus \mathcal{U}^{-1,n-1} \oplus \mathcal{U}^{1,n-1} \oplus \mathcal{U}^{1,n-3}.$$

But according to the hypothesis there is a trivialization ρ of $U^{0,n}$ such that $\bar{\partial}_{\mathcal{J}_1}\rho = 0$, so, in particular, $N_2 \cdot \rho$, the component of $\bar{\partial}_{\mathcal{J}_1}\rho$ in $\mathcal{U}^{-1,n-3}$, must vanish and since it vanishes on ρ , N_2 is the zero tensor. So we have in fact

$$d^H : \mathcal{U}^{0,n} \longrightarrow \mathcal{U}^{-1,n-1} \oplus \mathcal{U}^{1,n-1} \oplus \mathcal{U}^{1,n-3}.$$

and the last condition of Theorem (4.9) holds. \square

The same results hold for *negative SKT* structures. One should bear in mind, however that the relevant spaces for a negative structure are given by $W_-^k = \sum_{p-q=k} U^{p,q}$, that is, each W_-^k is an antidiagonal and integrability is equivalent to

$$d^H : \mathcal{W}_-^k \longrightarrow \mathcal{W}_-^{k+2} \oplus \mathcal{W}_-^k \oplus \mathcal{W}_-^{k-2},$$

or, in terms of the $U^{p,q}$ decomposition obtained by choosing a generalized complex extension, the only nontrivial component the Nijenhuis tensor is N_2 .

So an SKT structure (positive or negative) corresponds to a generalized Hermitian structure in which half of the Nijenhuis tensor of \mathcal{J}_2 vanishes.

5 Hodge theory

In this section we develop Hodge theory for manifolds with parallel Hermitian, bi-Hermitian and SKT structures. Our main result is that for a parallel positive Hermitian structure the Laplacian preserves the spaces $\mathcal{W}_+^{k,l}$ and hence induces a decomposition of the d^H -cohomology accordingly. This is in contrast with the fact that for usual manifolds the d^H -cohomology has only a \mathbb{Z}_2 -grading. For positive SKT manifolds, not only does the Laplacian preserve the spaces $\mathcal{W}_+^{k,l}$, but in fact there is an identity between the Laplacians for the operators d^H , δ_+^N and $\bar{\delta}_+^N$. We start with real Hodge theory and some operators of interest.

5.1 Differential operators, their adjoints and Laplacians

Given a generalized metric and orientation on a compact manifold M^m , we can form the Hodge star operator which gives us a positive definite inner product on forms:

$$\mathbb{G}(\varphi, \psi) = \int_M (\varphi, \star\psi)_{Ch} \quad \text{for all } \varphi, \psi \in \Omega^\bullet(M; \mathbb{R}). \quad (5.1)$$

Two basic results about d^H are:

Lemma 5.1. If M^m is compact

$$\int_M (d^H \varphi, \psi)_{Ch} = (-1)^m \int_M (\varphi, d^H \psi)_{Ch}. \quad (\text{Integration by parts})$$

$$\mathbb{G}(d^H \varphi, \psi) = \mathbb{G}(\varphi, (-1)^m \star^{-1} d^H \star \psi) \quad (\text{Formal adjoint})$$

hence the formal adjoint of d^H is given by $d^{H*} = (-1)^{\frac{m(m+1)}{2}} \star d^H \star$.

We will be mostly interested in the Dirac operators:

$$\begin{aligned} \mathcal{D}_+^H &= \frac{1}{2}(d^H - (-1)^{\frac{m(m-1)}{2}} \star d^H \star) = \frac{1}{2}(d^H + (-1)^{m+1} d^{H*}) \\ \mathcal{D}_-^H &= \frac{1}{2}(d^H + (-1)^{\frac{m(m-1)}{2}} \star d^H \star) = \frac{1}{2}(d^H + (-1)^m d^{H*}). \end{aligned} \quad (5.2)$$

The operators \mathcal{D}_{\pm}^H relate to the projections of d^H onto SD- and ASD- forms

$$\begin{aligned} d_{\pm}^H &: \Omega^{\bullet}(M) \longrightarrow \Omega_{\pm}^{\bullet}(M), \\ d_{\pm}^H \varphi &= \frac{1}{2}(1 \mp i^{-\frac{m(m-1)}{2}} \star) d^H \varphi. \end{aligned}$$

Lemma 5.2. Given $\varphi \in \Omega^{\bullet}(M)$ let φ_{\pm} be its SD and ASD components, then

$$\mathcal{D}_{\pm}^H \varphi = d_{\mp}^H \varphi_{+} + d_{\pm}^H \varphi_{-}.$$

In particular we see that \mathcal{D}_{-}^H preserves the spaces $\Omega_{\pm}^{\bullet}(M)$ while \mathcal{D}_{+}^H maps them to each other.

Proof.

$$\begin{aligned} \mathcal{D}_{\pm}^H \varphi &= \frac{1}{2}((d^H \mp (-1)^{\frac{m(m-1)}{2}} \star d^H \star)(\varphi_{+} + \varphi_{-})) \\ &= \frac{1}{2}((d^H \pm (-1)^{\frac{m(m-1)}{2}} i^{\frac{m(m-1)}{2}} \star d^H)\varphi_{+} + \frac{1}{2}((d^H \mp (-1)^{\frac{m(m-1)}{2}} i^{\frac{m(m-1)}{2}} \star d^H)\varphi_{-}) \\ &= \frac{1}{2}((d^H \pm i^{3\frac{m(m-1)}{2}} \star d^H)\varphi_{+} + \frac{1}{2}((d^H \mp i^{3\frac{m(m-1)}{2}} \star d^H)\varphi_{-}) \\ &= \frac{1}{2}((d^H \pm i^{-\frac{m(m-1)}{2}} \star d^H)\varphi_{+} + \frac{1}{2}((d^H \mp i^{-\frac{m(m-1)}{2}} \star d^H)\varphi_{-}) \\ &= d_{\mp}^H \varphi_{+} + d_{\pm}^H \varphi_{-}. \end{aligned}$$

□

Lemma 5.3. Let $\Delta^H = d^H d^{H*} + d^{H*} d^H$ be the d^H -Laplacian. Then

1. $\star \Delta^H = \Delta^H \star$ and
2. $(-1)^{m+1}(\mathcal{D}_{+}^H)^2 = (-1)^m(\mathcal{D}_{-}^H)^2 = \frac{1}{4}\Delta^H$.

Therefore Δ^H preserves the decomposition of forms into $\Omega_{\pm}^{\bullet}(M)$ and hence the d^H -cohomology of M splits as SD- and ASD-cohomology: $H_{d^H}(M) = \mathcal{H}_{+}^H(M) \oplus \mathcal{H}_{-}^H(M)$.

Not only does \mathcal{D}_{+}^H reverses the parity of forms but, according to Lemma 5.2, it swaps $\Omega_{+}^{\bullet}(M)$ with $\Omega_{-}^{\bullet}(M)$ and hence it is a generalization of the signature operator. Since \mathcal{D}_{+}^H has the same symbol as $d + (-1)^{m+1}d^*$ the indices of $\mathcal{D}_{+}^H : \Omega_{+}^{\bullet}(M) \longrightarrow \Omega_{-}^{\bullet}(M)$ and $\mathcal{D}_{+}^H : \Omega^{ev}(M) \longrightarrow \Omega^{od}(M)$ are just the usual signature and Euler characteristic, so we have

Lemma 5.4. The Euler characteristic, χ , and the signature, σ , of a compact manifold with closed 3-form (M, H) are given by

$$\chi = \dim(\mathcal{H}^{H,ev}(M)) - \dim(\mathcal{H}^{H,od}(M)); \quad \sigma = \dim(\mathcal{H}_{+}^H(M)) - \dim(\mathcal{H}_{-}^H(M)).$$

5.2 Signature and Euler characteristic of almost Hermitian manifolds

In a compact generalized almost Hermitian manifold we have sixteen operators induced by the $U^{p,q}$ decomposition of forms. It turns out that for these operators taking complex conjugates or adjoints are nearly the same thing. Firstly, we extend the real inner product on forms (5.1) to complex valued forms by requiring it to be Hermitian. If we let $\bar{\star}$ denote the operator given by $\bar{\star}\varphi = \star\bar{\varphi}$, we have

$$\mathbb{G}(\varphi, \psi) = (\varphi, \bar{\star}\psi) \quad \text{for all } \varphi, \psi \in \Omega^{\bullet}(M; \mathbb{C}).$$

Proposition 5.5. *Let M^{2n} be a compact generalized almost Hermitian manifold and let δ denote any of the operators $\delta_+, \bar{\delta}_+, \delta_-, \bar{\delta}_-, N_\alpha$ or \bar{N}_α , where $\alpha = 1, 2, 3, 4, \pm$. For $\varphi, \psi \in \Omega^\bullet(M; \mathbb{C})$ we have*

$$\int_M (\delta\varphi, \psi)_{Ch} = \int_M (\varphi, \delta\psi)_{Ch} \quad (\text{Integration by parts})$$

$$\delta^* = \pm \bar{\delta}, \quad (\text{Formal adjoint})$$

where the sign for the adjoint is positive if $\delta : \Omega_\pm(M; \mathbb{C}) \rightarrow \Omega_\pm(M; \mathbb{C})$ and negative otherwise.

Proof. The proof that integration by parts holds is the same for all of these operators, so we consider only δ_+ . It is enough to consider the case when $\varphi \in \mathcal{U}^{p,q}$ and hence $\delta_+\varphi \in \mathcal{U}^{p+1,q+1}$ and it pairs trivially with $\mathcal{U}^{k,l}$, unless $k = -p - 1$ and $l = -q - 1$. Hence we may assume that $\psi \in \mathcal{U}^{-p-1,-q-1}$ and compute

$$\int_M (\delta_+\varphi, \psi)_{Ch} = \int_M (d^H\varphi, \psi)_{Ch} = \int_M (\varphi, d^H\psi)_{Ch} = \int_M (\varphi, \delta_+\psi)_{Ch},$$

where in the first equality we have used that the remaining components of $d^H\varphi$ do not lie in $\mathcal{U}^{p+1,q+1}$, hence they pair trivially with ψ , in the second equality we integrated by parts and then reversed the argument.

For the formal adjoint, again taking $\delta = \delta_+$, $\varphi \in \mathcal{U}^{p,q}$ and $\psi \in \mathcal{U}^{p+1,q+1}$ we compute:

$$\begin{aligned} \mathbb{G}(\delta_+\varphi, \psi) &= \int_M (\delta_+\varphi, \bar{\star}\psi)_{Ch} = i^{-p-q-2} \int_M (\varphi, \delta_+\bar{\psi})_{Ch} \\ &= i^{-p-q-2} \int_M (\varphi, (-1)^n \bar{\star} \bar{\delta}_+\bar{\psi})_{Ch} = i^{-p-q-2} (-1)^n \int_M (\varphi, \bar{\star} \star \bar{\delta}_+\psi)_{Ch} \\ &= i^{-2p-2q-2} (-1)^n \int_M (\varphi, \bar{\star} \bar{\delta}_+\psi)_{Ch} = -\mathbb{G}(\varphi, \bar{\delta}_+\psi), \end{aligned}$$

where we have used several times that on $\mathcal{U}^{p,q}$ \star is multiplication by i^{p+q} and in the last equality we used that $p + q = n \pmod{2}$. \square

Therefore we can form the Dirac operator corresponding to, say, δ_+ :

$$\not\delta_+ = \delta_+ - \delta_+^* = \delta_+ + \bar{\delta}_+;$$

Since the symbol sequence of δ_+ is exact, we get elliptic operators for the sequences (2.9) and (2.10):

$$\begin{aligned} \not\delta_+ : \Omega^{ev}(M) &\rightarrow \Omega^{od}(M); \\ \not\delta_+ : \Omega_+(M) &\rightarrow \Omega_-(M); \end{aligned} \quad (5.3)$$

The indices of these operators are just the Euler characteristic and the signature of M :

Theorem 5.6. *Let (M, H) be a compact manifold with a 3-form H and let $(\mathcal{G}, \mathcal{J}_1)$ be an almost generalized Hermitian structure on M . Let*

$$h^{ev/od} = \dim(\ker(\not\delta_+ : \Omega^{ev/od} \rightarrow \Omega^{od/ev})),$$

$$w^\pm = \dim(\ker(\not\delta_\pm : \Omega_\pm(M) \rightarrow \Omega_\mp(M)));$$

Then

$$\chi(M) = h^{ev} - h^{od} \quad \text{and} \quad \sigma(M) = w^+ - w^-.$$

Proof. To prove the claim about the Euler characteristic we only have to show that the symbols of δ_+ and $\mathcal{D}_+^H : \Omega^{ev} \rightarrow \Omega^{od}$ since due to Lemma 5.4 the index of \mathcal{D}_+^H is the Euler characteristic. To compare the symbols we have

$$\mathcal{D}_+^H = d^H - (d^H)^* \sim \delta_+ + \bar{\delta}_+ + \delta_- + \bar{\delta}_- - (-\bar{\delta}_+ - \delta_+ + \bar{\delta}_- + \delta_-) = 2\delta_+ + 2\bar{\delta}_+ = 2\delta_+, \quad (5.4)$$

where \sim means that the operators have the same symbol and in the second passage we used Proposition 5.5.

The proof of the claim regarding the signature is done in the same lines. \square

5.3 Hodge theory on parallel Hermitian manifolds

Theorem 5.7. *Let (M, H) be a compact manifold with 3-form, let $(\mathcal{G}, \mathcal{I}_+, \mathcal{I}_-)$ be a generalized metric with a pair of almost Hermitian structures and $\mathcal{J}_1 = \mathcal{I}_+ + \mathcal{I}_-$. Then*

1. *If \mathcal{I}_+ (resp. \mathcal{I}_-) is a parallel positive (resp. negative) Hermitian structure, the d^H -Laplacian preserves the spaces $\mathcal{W}_+^{k,l}$ (resp. $\mathcal{W}_-^{k,l}$) and hence the d^H -cohomology of M inherits a corresponding $\mathbb{Z} \times \mathbb{Z}_2$ -grading;*
2. *If $(\mathcal{G}, \mathcal{J}_1)$ is a parallel bi-Hermitian structure, the d^H -Laplacian preserves the spaces $\mathcal{U}^{p,q}$ and hence the d^H -cohomology of M inherits a corresponding \mathbb{Z}^2 -grading;*

Proof. The claim for positive and negative structures are analogous and together they imply the last claim, so it is enough to prove the first claim.

If (M, H) has a generalized Hermitian structure $(\mathcal{G}, \mathcal{J}_1)$, then, since each diagonal of the $\mathcal{U}^{p,q}$ decomposition lies in either $\Omega_+^\bullet(M; \mathbb{C})$ or in $\Omega_-^\bullet(M; \mathbb{C})$, so Lemma 5.2 implies that \mathcal{D}_+^H and \mathcal{D}_-^H are decomposed into a sum of operators:

$$\begin{aligned} \mathcal{D}_+^H &= \delta_+ + N_1 + N_4 + N_+ + \bar{\delta}_+ + \bar{N}_1 + \bar{N}_4 + \bar{N}_+; \\ \mathcal{D}_-^H &= \delta_- + N_2 + N_3 + N_- + \bar{\delta}_- + \bar{N}_2 + \bar{N}_3 + \bar{N}_-. \end{aligned} \quad (5.5)$$

If \mathcal{I}_+ is parallel, then, according to Theorem 3.4, $N_2 = N_3 = 0$ and from (5.5) we have that $\mathcal{D}_-^H = \delta_- + \bar{\delta}_- + N_- + \bar{N}_-$, which clearly preserves the \mathcal{W}_+^k decomposition, therefore $\Delta^H = 4(\mathcal{D}_-^H)^2$ also preserves the \mathcal{W}^k decomposition of forms. Since the Laplacian has even parity, it also preserves $\Omega^{ev}(M)$ and $\Omega^{od}(M)$, hence the Laplacian preserves the spaces $\mathcal{W}_+^{k,l}$. \square

Corollary 5.8. Let M be a manifold with 3-form.

- An SKT structure on M induces a $\mathbb{Z} \times \mathbb{Z}_2$ -grading on the d^H -cohomology;
- (Gualtieri [19]) A generalized Kähler structure induces a \mathbb{Z}^2 -grading on the d^H -cohomology.

For each of the cases covered in the previous theorem, we denote by $H_{d^H}^{\bullet,\bullet}(M)$ the corresponding decomposition of the d^H -cohomology.

Corollary 5.9. (Riemann bilinear relations) In a compact parallel positive Hermitian manifold

$$i^{-k}(a, \bar{a})_{Ch} > 0 \quad \text{for all } a \in H_{d^H}^{k,l}(M) \setminus \{0\}.$$

Proof. Indeed, let α be the harmonic representative for the class $a \in H_{d^H}^{k,l}(M) \setminus \{0\}$. Since $\alpha \in \mathcal{W}_+^{k,l}$, $\star\alpha = i^k\alpha$ and hence $\bar{\star}\alpha = i^{-k}\bar{\alpha}$ and

$$i^{-k}(\alpha, \bar{\alpha})_{Ch} = (\alpha, \bar{\star}\alpha)_{Ch} > 0.$$

\square

Finally in this context we get a clearer version of Theorem 5.6:

Corollary 5.10. Letting $w^{k,l}$ be the dimension of the space of harmonic sections of $\mathcal{W}_+^{k,l}$ in a parallel Hermitian manifold we have

$$\chi(M) = \sum (-1)^l w^{k,l} \quad \text{and} \quad \sigma(M) = \sum (-1)^{\frac{n-k}{2}} w^{k,l}.$$

5.4 Hodge theory on SKT manifolds

While for parallel Hermitian manifolds we could prove that the Laplacian preserves the spaces $\mathcal{W}_+^{k,l}$, in and SKT manifold we can go further and show that there is an identity of Laplacians:

Theorem 5.11. *In a positive SKT manifold,*

$$\Delta_{\delta_+^N} = \Delta_{\overline{\delta_+^N}} = \frac{1}{4} \Delta^H.$$

Proof. From $\mathcal{D}_+^H = \delta_+^N + \overline{\delta_+^N}$, Lemma 5.3 and Proposition 5.5 we have

$$\frac{1}{4} \Delta^H = -(\mathcal{D}_+^H)^2 = -(\delta_+^N + \overline{\delta_+^N})^2 = -(\delta_+^N - \delta_+^{N*})^2 = \Delta_{\delta_+^N}$$

□

5.5 Relation to Dolbeault cohomology

As we saw in Proposition 4.1, given a positive SKT structure $(\mathcal{G}, \mathcal{I})$ on M one can extend \mathcal{I}_+ to a generalized complex structure \mathcal{J}_1 of complex type. In this case, in the metric splitting of $\mathbb{T}M$, the structures \mathcal{J}_1 and $\mathcal{J}_2 = \mathcal{G}\mathcal{J}_1$ are given by

$$\mathcal{J}_1 = \begin{pmatrix} I & 0 \\ 0 & -I^* \end{pmatrix}, \quad \mathcal{J}_2 = \begin{pmatrix} 0 & -\omega^{-1} \\ \omega & 0 \end{pmatrix},$$

where $\omega = g \circ I$ and for such generalized Hermitian structure Proposition 1.8 gives an automorphism of the space of forms which relates the $U^{p,q}$ decomposition of forms with the usual $\wedge^{p,q} T^*M$ decomposition of forms determined by the complex structure I .

In this section we relate these two decomposition of forms and corresponding cohomologies. Precisely, the effect of applying the automorphism Ψ from Proposition 1.8 to $\wedge^\bullet T_{\mathbb{C}}^*M$ on d^H is simply to conjugate it by Ψ , so we get a new operator

$$\hat{d}^H = \Psi^{-1} \circ d^H \circ \Psi.$$

Since d^H splits according to the $U^{p,q}$ into six operators, the same is true for \hat{d}^H and we can define

$$\begin{aligned} \hat{\delta}_+ &: \Omega^{p,q} \longrightarrow \Omega^{p-1,q}; & \hat{\delta}_+^{\wedge} &: \Omega^{p,q} \longrightarrow \Omega^{p+1,q}, \\ \hat{\delta}_- &: \Omega^{p,q} \longrightarrow \Omega^{p,q+1}; & \hat{\delta}_-^{\wedge} &: \Omega^{p,q} \longrightarrow \Omega^{p,q+1}, \\ \hat{N} &: \Omega^{p,q} \longrightarrow \Omega^{p-1,q-2} & \hat{N}^{\wedge} &: \Omega^{p,q} \longrightarrow \Omega^{p+1,q+2}; \end{aligned}$$

$$\hat{d}^H = \hat{\delta}_+ + \hat{\delta}_+^{\wedge} + \hat{\delta}_- + \hat{\delta}_-^{\wedge} + \hat{N} + \hat{N}^{\wedge}.$$

Proposition 5.12. *Let $(\mathcal{G}, \mathcal{I}_+)$ be a positive SKT structure on a manifold with 3-form (M, H) , let \mathcal{J} be the generalized complex extension of \mathcal{I}_+ given by Proposition 4.1. Let*

$$\partial^{\omega^{-1}} = \{\partial, \omega^{-1}\} \quad \text{and} \quad \bar{\partial}^{\omega^{-1}} = \{\bar{\partial}, \omega^{-1}\},$$

let ζ_i be the component of $e_*^{\frac{i}{2}\omega^{-1}}\bar{\partial}\omega$ lying in $\wedge^i T \otimes \wedge^{3-i} T^*$ and let χ be the $(1,2)$ -component of $[\omega^{-1}, \omega^{-1}]_{SN}$, where $[\cdot, \cdot]_{SN}$ is the Schouten–Nijenhuis bracket. Then the following hold

$$\begin{aligned}\hat{\delta}_+ &= \frac{i}{2}\bar{\partial}\omega^{-1} + 2i\zeta_2 & \hat{\delta}_+ &= \partial \\ \hat{\delta}_- &= \bar{\partial} + 2i\zeta_1 & \hat{\delta}_- &= \frac{i}{2}\partial\omega^{-1} \\ \hat{N} &= \frac{1}{8}\chi & \hat{N} &= 2i\bar{\partial}\omega.\end{aligned}\tag{5.6}$$

Proof. The proof is a direct computation of the operator \hat{d}^H :

$$\begin{aligned}\hat{d}^H &= e^{-\frac{i}{2}\omega^{-1}}e^{-i\omega}(d + H\wedge)e^{i\omega}e^{\frac{i}{2}\omega^{-1}} \\ &= e^{-\frac{i}{2}\omega^{-1}}e^{-i\omega}e^{i\omega}(d + (H + id\omega)\wedge)e^{\frac{i}{2}\omega^{-1}} \\ &= e^{-\frac{i}{2}\omega^{-1}}de^{\frac{i}{2}\omega^{-1}} + 2ie^{-\frac{i}{2}\omega^{-1}}(\bar{\partial}\omega\wedge)e^{\frac{i}{2}\omega^{-1}}\end{aligned}\tag{5.7}$$

We compute separately each of the two operators above making up \hat{d}^H :

$$e^{-\frac{i}{2}\omega^{-1}}de^{\frac{i}{2}\omega^{-1}} = d + [d, \frac{i}{2}\omega^{-1}] + \frac{1}{2!}[[d, \frac{i}{2}\omega^{-1}], \frac{i}{2}\omega^{-1}] + \frac{1}{3!}[[[d, \frac{i}{2}\omega^{-1}], \frac{i}{2}\omega^{-1}], \frac{i}{2}\omega^{-1}] + \dots$$

The first term is just $d = \partial + \bar{\partial}$, while the second term is a version of the symplectic adjoint of d now obtained in a nonintegrable setting, $\delta\omega^{-1} = d \circ \omega^{-1} - \omega^{-1} \circ d$. Since ω is not closed, $\delta\omega^{-1}$ does not square to zero. Integrability of the complex structure gives $d = \partial + \bar{\partial}$ and recalling that ω^{-1} is of type $(1,1)$ we have

$$\delta\omega^{-1} = \partial\omega^{-1} + \bar{\partial}\omega^{-1}; \partial\omega^{-1} : \Omega^{p,q} \longrightarrow \Omega^{p,q-1}; \quad \bar{\partial}\omega^{-1} : \Omega^{p,q} \longrightarrow \Omega^{p-1,q}.$$

Since ω^{-1} is even, the third term in the series coincides with a multiple of the expression for the derived bracket of ω^{-1} with itself, that is, the Schouten–Nijenhuis bracket:

$$[[d, \frac{i}{2}\omega^{-1}], \frac{i}{2}\omega^{-1}] = -\{\{\frac{i}{2}\omega^{-1}, d\}, \frac{i}{2}\omega^{-1}\} = \frac{1}{4}[\omega^{-1}, \omega^{-1}]_{SN}.$$

Since ω^{-1} is of type $(1,1)$, integrability of the complex structure implies that this term this is a 3-vector lies in $\wedge^{2,1}T \oplus \wedge^{1,2}T$, so this term decomposes as

$$[\omega^{-1}, \omega^{-1}]_{SN} = \chi + \bar{\chi}, \quad \chi \in \wedge^{1,2}TM.$$

The fourth term is the commutator $[[\omega^{-1}, \omega^{-1}]_{SN}, \omega^{-1}]$ which vanishes since ω^{-1} is a bivector and hence so do the remaining terms in the series. So we have established that

$$e^{-\frac{i}{2}\omega^{-1}}de^{\frac{i}{2}\omega^{-1}} = \partial + \bar{\partial} + \frac{i}{2}\partial\omega^{-1} + \frac{i}{2}\bar{\partial}\omega^{-1} + \frac{1}{8}(\chi + \bar{\chi})\tag{5.8}$$

Next we compute the second summand in (5.7):

$$\begin{aligned}2ie^{-\frac{i}{2}\omega^{-1}} \circ \bar{\partial}\omega \circ e^{\frac{i}{2}\omega^{-1}} &= 2ie^{-\frac{i}{2}\omega^{-1}} \circ (e_*^{\frac{i}{2}\omega^{-1}} e_*^{-\frac{i}{2}\omega^{-1}} \bar{\partial}\omega) \circ e^{\frac{i}{2}\omega^{-1}} \\ &= 2ie^{-\frac{i}{2}\omega^{-1}} \circ e^{\frac{i}{2}\omega^{-1}} \circ (e_*^{-\frac{i}{2}\omega^{-1}} \bar{\partial}\omega) \\ &= 2ie_*^{-\frac{i}{2}\omega^{-1}} \bar{\partial}\omega,\end{aligned}$$

where we have used (1.1) in the second equality. The element $e_*^{-\frac{i}{2}\omega^{-1}}\bar{\partial}\omega \in \wedge^3\mathbb{T}_{\mathbb{C}}M$ has six components:

$$e_*^{-\frac{i}{2}\omega^{-1}}\bar{\partial}\omega \in (\wedge^{1,2}T^*) \oplus (T^{1,0} \otimes \wedge^{1,1}T^*) \oplus (T^{0,1} \otimes \wedge^{0,2}T^*) \oplus (\wedge^{2,0}T \otimes T^{*1,0}) \oplus (\wedge^{1,1}T \otimes T^{*0,1}) \oplus (\wedge^{2,1}T)$$

Since, for a 1-form ξ , $e_*^{-\frac{i}{2}\omega^{-1}}\xi = \frac{i}{2}\omega^{-1}(\xi) + \xi$, we see that the 3-form component is $\bar{\partial}\omega$ and that the 3-vector component is $-\frac{i}{8}\omega^{-1}\bar{\partial}\omega$. We let ζ_1 be the component of $e_*^{-\frac{i}{2}\omega^{-1}}\bar{\partial}\omega$ in $T \otimes \wedge^2 T^*$ and ζ_2 be the component in $\wedge^2 T \otimes T^*$, so that $\zeta_1 : \wedge^{p,q} \rightarrow \wedge^{p,q+1}$, $\zeta_2 : \wedge^{p,q} \rightarrow \wedge^{p-1,q}$, and

$$2ie^{-\frac{i}{2}\omega^{-1}} \circ \bar{\partial}\omega \circ e^{\frac{i}{2}\omega^{-1}} = 2i\bar{\partial}\omega + 2i\zeta_1 + 2i\zeta_2 + \frac{1}{4}\omega^{-1}(\bar{\partial}\omega)$$

Putting this together with (5.8) and the fact that d^H does not have a component mapping $\Omega^{p,q}(M)$ to $\Omega^{p-2,q-1}(M)$ we conclude that $\bar{\chi} = -2\omega^{-1}(\bar{\partial}\omega)$ and obtain (5.6). \square

Next, we let $\partial^{i\bar{\partial}\omega}$ be the operator $\partial + i\bar{\partial}\omega \wedge$. Then we observe that even though $\partial^{i\bar{\partial}\omega}$ does not preserve the degree of a form, it preserves the holomorphic degree and the parity of the anti-holomorphic degree, i.e.,

$$\partial^{i\bar{\partial}\omega} : \Omega^{p,q}(M) \rightarrow \Omega^{p+1,q}(M), \quad p \in \mathbb{Z}, \quad q \in \mathbb{Z}_2.$$

Therefore its cohomology has a natural $\mathbb{Z} \times \mathbb{Z}_2$ -grading. For an operator δ with $\delta^2 = 0$ we denote its cohomology by $H_\delta(M)$.

Corollary 5.13. Let $(\mathcal{G}, \mathcal{I}_+)$ be a positive SKT structure on a compact manifold, let \mathcal{J} be the generalized complex extension from Proposition 4.1 and let (g, I) be the corresponding Hermitian structure with Hermitian form $\omega = g \circ I$. Then for $p, q \in \mathbb{Z}$

$$H_\partial^{p,q}(M) \cong H_{\delta_+}^{q-p, n-p-q}(M),$$

and

$$H_{\partial^{i\bar{\partial}\omega}}^p(M) \cong H_{\delta_+^N}^{n-2p}(M) \cong H_{d^H}^{n-2p}(M).$$

Proof. Indeed, Ψ puts $\Omega^{p,q}(M)$ in correspondence with $\mathcal{U}^{q-p, n-p-q}$ and $\Omega^{p,\bullet}(M)$ with \mathcal{W}^{n-2p} and according to Proposition 5.12, $\Psi \circ \partial = \bar{\delta}_+ \circ \Psi$ and $\Psi \circ \partial^{2i\omega} = \bar{\delta}_+^N \circ \Psi$. Therefore $H_\partial^{p,q}(M) \cong H_{\delta_+}^{q-p, n-p-q}(M)$ and $H_{\partial^{i\bar{\partial}\omega}}^p(M) \cong H_{\delta_+^N}^{n-2p}(M)$. Of course the cohomology of $\partial^{2i\bar{\partial}\omega}$ is the same as the cohomology if $\partial^{i\bar{\partial}\omega}$, as the automorphism

$$m : \wedge^{\bullet,\bullet} T^* M \rightarrow m : \wedge^{\bullet,\bullet} T^* M \quad m(\alpha) = 2^{\frac{q}{2}} \alpha \quad \text{for all } \alpha \in \wedge^{p,q} T^* M,$$

relates them. \square

Interestingly we have that $d^H = \partial^{i\bar{\partial}\omega} + \bar{\partial}^{-i\partial\omega}$ and the operators $\partial^{i\bar{\partial}\omega}$ and $\bar{\partial}^{-i\partial\omega}$ both square to zero (due to the SKT condition $\partial\bar{\partial}\omega = 0$) and graded commute. Yet, they are not well behaved with respect to the \mathbb{Z}^2 -bigrading of forms and only on a \mathbb{Z}_2^2 -grading do they behave nicely, as there we have

$$\partial^{i\bar{\partial}\omega} : \Omega^{p,q}(M) \rightarrow \Omega^{p+1,q}(M) \quad \bar{\partial}^{-i\partial\omega} : \Omega^{p,q}(M) \rightarrow \Omega^{p,q+1}(M), \quad (p, q) \in \mathbb{Z}_2^2.$$

There is a standard trick to recover one \mathbb{Z} -grading, say the ‘ p ’-grading. We introduce a new formal variable, β , alter the operators $\partial^{i\bar{\partial}\omega}$ and $\bar{\partial}^{-i\partial\omega}$ so that they act on $E_1 = \Omega^{p,q}(M) \otimes \langle \beta \rangle$:

$$\partial^{i\bar{\partial}\omega}(\alpha\beta^k) = (\partial^{i\bar{\partial}\omega}\alpha)\beta^k \quad \bar{\partial}^{-i\partial\omega}(\alpha\beta^k) = (\bar{\partial}\alpha)\beta^k - i\partial\omega \wedge \alpha\beta^{k+1}$$

and introduce a $\mathbb{Z} \times \mathbb{Z}_2$ -grading on E_1 by declaring that for $\Omega^{p,q}(M) \otimes \beta^k = E_1^{p-2k,q}$, $(p-2k, q) \in \mathbb{Z} \times \mathbb{Z}_2$. I.e., β has degree $(-2, 0)$. With this arrangement

$$\partial^{i\bar{\partial}\omega} : E_1^{p,q}(M) \rightarrow E_1^{p+1,q}(M) \quad \bar{\partial}^{-i\partial\omega} : E_1^{p,q}(M) \rightarrow E_1^{p,q+1}(M), \quad (p, q) \in \mathbb{Z} \times \mathbb{Z}_2.$$

In this setting, one can produce a $\mathbb{Z} \times \mathbb{Z}_2$ -graded spectral sequence whose first page is $H_{\partial^{i\bar{\partial}\omega}}^\bullet(M) \otimes \langle \beta \rangle$ and the last is $H_{d^H}^\bullet(M) \otimes \langle \beta \rangle$. Since these are isomorphic, by Corollary 5.13, this sequence degenerates at the first page.

Theorem 5.14. (SKT spectral sequence) *In an SKT manifold the decomposition $d^H = (\partial + i\bar{\partial}\omega) + (\bar{\partial} - i\partial\omega)$ gives rise to a spectral sequence which degenerates at the first page.*

On a more general note, the isomorphism Ψ used to prove Proposition 5.12 can be used even in the nonintegrable case to give us information about the Euler characteristic and signature of almost Hermitian manifolds. Indeed, according to Proposition 1.8 we have

$$\Psi : \bigoplus_q \wedge^{p,q} T^*M \cap \Omega^{ev}(M) \longrightarrow W_+^{k,0}; \quad \Psi : \bigoplus_q \wedge^{p,q} T^*M \cap \Omega^{od}(M) \longrightarrow W_+^{k,1}.$$

And hence conjugating $\bar{\delta}_+$ by Ψ we get an operator

$$\hat{\delta}_+ = \Psi \circ \bar{\delta}_+ \circ \Psi^{-1} : \Omega^{p,q} \longrightarrow \Omega^{p+1,q}.$$

Since the isomorphism Ψ from Proposition 1.8 identifies $T^{*1,0}M$ with $V_+^{1,0}$, one can readily check that ∂ and $\hat{\delta}_+$ have the same symbol and hence the same index. Of course complex conjugation swaps ∂ and $\bar{\partial}$, allowing us to translate Theorem 5.6 to Dolbeault cohomology terms:

Corollary 5.15. Let (M, I, g) be a compact almost Hermitian manifold, $\bar{\partial}^* = \bar{\partial} - \bar{\partial}^*$ and, for $q, k \in \mathbb{Z}_2$,

$$\begin{aligned} w^q &= \dim(\ker \bar{\partial}^* : \bigoplus_p \Omega^{p,q}(M) \longrightarrow \bigoplus_p \Omega^{p,q+1}(M)) \\ h^k &= \dim(\ker \bar{\partial}^* : \Omega^k(M) \longrightarrow \Omega^{k+1}(M)) \end{aligned}$$

Then

$$\chi(M) = h^{ev} - h^{od} \quad \text{and} \quad \sigma(M) = w^{ev} - w^{od}.$$

In particular in a compact complex manifold if $h^{p,q} = \dim(H_{\bar{\partial}}^{p,q}(M))$, then

$$\chi = \sum_{p,q} (-1)^{p+q} h^{p,q} \quad \text{and} \quad \sigma = \sum_{p,q} (-1)^q h^{p,q}(M). \quad (5.9)$$

Remark. Using Frölicher's spectral sequence, Frölicher proved the identity for the Euler characteristic assuming integrability of the complex structure [14]. The identity regarding the signature is an extension of the Hodge Index Theorem, which, in its modern version (see, e.g., [32], Theorem 6.33), states that (5.9) holds on a compact Kähler manifold and is obtained as a consequence of the development of Hodge theory of Kähler manifolds. The main point of Corollary 5.15 is that these identities do not depend on existence of Kähler structures or on the integrability of the complex structure, so, in effect, *they do not represent an obstruction to the existence of any of the structures studied here.* This is to be compared with Theorem 5.14 which provides the first nontrivial differential-topological obstruction to the existence of SKT structures on complex manifolds known to the author.

Example 5.16 (Calabi–Eckman manifolds). In this example we show that the Calabi–Eckman manifolds $M_{u,v} \cong S^{2u+1} \times S^{2v+1}$ do not admit SKT structures for $v \geq u > 1$, the reason being that the d^H -cohomology of these manifolds can not be isomorphic to the $\bar{\partial}^{i\bar{\partial}\omega}$ -cohomology, and hence Corollary 5.13 fails for these manifolds.

Indeed, if such a manifold had an SKT structure, then $[d^c\omega] = 0$ and the d^H -cohomology would be isomorphic to the de Rham cohomology. On the other hand, the Dolbeault cohomology of these manifolds was computed by Borel [22] and is given by

$$H_{\bar{\partial}}^{\bullet,\bullet}(M_{u,v}) = \wedge \text{span}\{x^{0,1}, z^{v+1,v}\} \otimes \mathbb{C}[y^{1,1}]/y^{u+1},$$

where $a^{p,q}$ is a generator of bidegree (p, q) . From this, one immediately sees that $H_{\bar{\partial}}^{2,1}(M_{u,v}) = \{0\}$ and hence $\partial\omega$ would be the trivial Dolbeault class and the $\bar{\partial}^{i\partial\omega}$ -cohomology would be isomorphic to the usual Dolbeault cohomology. Since the Dolbeault cohomology of the Calabi-Eckman manifolds is not isomorphic to the de Rham cohomology, $M_{u,v}$ can not be SKT. \blacksquare

5.6 Hermitian symplectic structures

A particular type of SKT structure for which Theorem 5.14 is particularly relevant are the so called *Hermitian symplectic structures*. These consist of a pair (I, ω) of integrable complex structure and symplectic structure such that $\omega(X, IX)$ is positive for every nonzero vector X . The difference between these structures and Kähler structures is that here we do not require ω to be of type $(1,1)$. Yet, we can decompose ω into its (p, q) components with respect to the complex structure and then one readily obtains that

$$d^c\omega^{1,1} = id(\omega^{2,0} - \omega^{0,2})$$

hence $(I, \omega^{1,1})$ is an SKT structure and $H = id(\omega^{2,0} - \omega^{0,2})$ is exact hence the d^H -cohomology is isomorphic to the de Rham cohomology and $H^{1,2} = i\partial\omega^{0,2}$ is ∂ -exact hence the $\bar{\partial}^{-i\bar{\partial}\omega^{1,1}}$ -cohomology is isomorphic to the ∂ -cohomology and Theorem 5.14 gives:

Corollary 5.17. In a compact Hermitian symplectic manifold the Frölicher spectral sequence degenerates at the second page, i.e., the Dolbeault and the de Rham cohomologies are isomorphic as graded vector spaces.

6 Hodge theory beyond $U(n)$

The decomposition of harmonics into their (p, q) -components in a Kähler manifold is a phenomenon that repeats itself for any other special holonomy group and, as such, is a result on Riemannian geometry. This approach is quite different from what we have done so far as, just like in the original Kähler identities, we relied on the underlying complex structures heavily to develop our theory. This section we show that with the appropriate setup, our results can also be extended to any other holonomy groups. Throughout this section we let $\{e_1, \dots, e_m\} \in \Gamma(TM)$ be an orthonormal frame and $\{e^1, \dots, e^m\} \in \Gamma(T^*M)$ be its dual frame.

As before, given an oriented Riemannian manifold with closed 3-form, (M, g, H) we consider the metric connection $\hat{\nabla}^\pm$ whose torsion is $\mp H$ and let ∇ denote the Levi-Civita connection. Using the orthonormal frame, we can write explicit expressions for ∇^\pm . Indeed, if we define $h_{ijk} = H(e_i, e_j, e_k)$, then, using Einstein summation convention and omitting the symbol for the wedge product, H is given by

$$H = \frac{1}{3!} h_{ijk} e^i e^j e^k.$$

and we have

$$\nabla^\pm = \nabla \mp \frac{1}{4} h_{ijk} e^i e^j e_k. \quad (6.1)$$

If the holonomy of ∇^+ is the Lie group G_+ , then using the isomorphism $TM \cong V_+$, we realize its Lie algebra, \mathfrak{g}_+ , as a sub Lie algebra of $\mathfrak{so}(V_+) = \wedge^2 V_+$. *Mutatis mutandis*, the same holds for ∇^- and we get $\mathfrak{g}_- \subset \wedge^2 V_-$. Next we notice that $\mathfrak{g}_+ \oplus \mathfrak{g}_- \subset \wedge^2 TM = \mathfrak{spin}(TM)$ and hence the elements of $\mathfrak{g}_+ \oplus \mathfrak{g}_-$ act on forms, thought of as spinors. Further, since V_+ is orthogonal to V_- and $\mathfrak{g}_\pm \subset \wedge^2 V_\pm$, the Lie algebra action of \mathfrak{g}_+ and \mathfrak{g}_- on forms commute and we get an action of $\mathfrak{g}_+ \oplus \mathfrak{g}_-$ on forms as a direct sum of the individual actions of the Lie algebras. Now we are in condition to state the main theorem of this section, which extends Theorem 5.7 to general holonomy groups.

Theorem 6.1. *Let (M, g, H) be a compact Riemannian manifold endowed with a closed 3-form. If ∇^\pm , the metric connections with torsion $\mp H$, have holonomy in G_\pm , then the d^H -cohomology of*

M splits according to the decomposition of forms into irreducible representations of the action of $\mathfrak{g}_+ \oplus \mathfrak{g}_- \subset \wedge^2 V_+ \oplus \wedge^2 V_-$ on forms.

Lemma 6.2. The connections ∇^\pm preserves the irreducible representations of \mathfrak{g}_\pm , respectively.

Proof. Indeed, if we denote by $\widetilde{\mathfrak{g}}_+ \subset \mathfrak{spin}(TM) \cong \wedge^2 T^*M$ the bundle of endomorphisms of TM defined by the connection ∇^+ , the condition that ∇^+ has reduced holonomy implies that this bundle is preserved by parallel transport. Now \mathfrak{g}_+ is simply the image of $\widetilde{\mathfrak{g}}_+$ by the parallel isomorphism $\text{Id} + g : T^*M \rightarrow V_+$, hence the bundle $\mathfrak{g}_+ \subset \wedge^2 V_+$ is also parallel and its irreducible representations are preserved by the connection. \square

Next we define

$$\varepsilon_\pm^i = e^i \pm e_i \in \Gamma(V_\pm)$$

and

$$H_\pm = \frac{1}{3!} h_{ijk} \varepsilon_\pm^i \varepsilon_\pm^j \varepsilon_\pm^k \in \text{Clif}^3(V_\pm) \subset \text{Clif}^3(TM).$$

To prove the theorem we need to extend to the torsion case the formulas relating the Levi–Civita connection with the exterior derivative and its adjoint:

$$d = e^i \wedge \nabla_{e_i}; \quad d^* = -\iota_{e_i} \nabla_{e_i}. \quad (6.2)$$

To avoid considering separate cases according to whether the dimension of the manifold is even or odd, we let

$$\mathcal{D}_+ = d^H - d^{H*} \quad \text{and} \quad \mathcal{D}_- = d^H + d^{H*}.$$

Lemma 6.3. With the definitions above

$$\mathcal{D}_+^H = \varepsilon_+^i \nabla_{e_i}^+ + H_+; \quad 2\mathcal{D}_-^H = \varepsilon_-^i \nabla_{e_i}^- + H_-. \quad (6.3)$$

Proof. Since the statements are similar, we will only prove the first identity. The left hand side of (6.3) is

$$\begin{aligned} \mathcal{D}_+^H &= d + H - d^* - H^* \\ &= d - d^* + \frac{1}{3!} h_{ijk} e^i e^j e^k + \frac{1}{3!} h_{ijk} e_i e_j e_k. \end{aligned} \quad (6.4)$$

Next, we use relation (6.1) to re-write the right hand side as

$$\varepsilon_+^i \nabla_{e_i}^+ + H_+ = e^i \wedge \nabla_{e_i} + \iota_{e_i} \nabla_{e_i} - \frac{1}{2} h_{ijk} e^i e^j e_k - \frac{1}{2} h_{ijk} e_i e_j e_k + H_+.$$

Due to (6.2), the first two terms are $d - d^*$. Expanding H_+ we get

$$\varepsilon_+^i \nabla_{e_i}^+ + H_+ = d - d^* - \frac{1}{2} h_{ijk} e^i e^j e_k - \frac{1}{2} h_{ijk} e_i e_j e_k + \frac{1}{3!} h_{ijk} e^i e^j e^k + \frac{1}{2!} h_{ijk} e^i e^j e_k + \frac{1}{2!} h_{ijk} e^i e_j e_k + \frac{1}{3!} h_{ijk} e_i e_j e_k.$$

which equals (6.4). \square

Proof of Theorem 6.1. Since ∇^+ preserves the irreducible representations of \mathfrak{g}_+ , Lemma 6.3 implies that \mathcal{D}_+^H preserves the irreducible representations of $\mathfrak{g}_+ \oplus \text{Clif}(V_-)$ and hence so does the d^H -Laplacian. Similarly, \mathcal{D}_-^H preserves the irreducible representations of $\text{Clif}(V_+) \oplus \mathfrak{g}_-$ and hence Δ_{d^H} preserves the intersections of these representations, which are just the irreducible representations of $\mathfrak{g}_+ \oplus \mathfrak{g}_-$. \square

6.1 Integrability

A simple consequence of (6.2) is that if $W \subset \wedge^k T^*M$ is a representation of the holonomy group of the Levi-Civita connection, then the exterior derivative restricted to sections of W can only land in representations present in $T^*M \wedge W$. This is the Riemannian version of the claim that in a complex manifold $d : \Omega^{p,q}(M) \rightarrow \Omega^{p+1,q}(M) \oplus \Omega^{p,q+1}(M)$. Lemma 6.3 has similar implications.

Proposition 6.4. *Let (M, g, H) be a compact oriented Riemannian manifold with a closed 3-form and let $\mathfrak{g}_\pm \subset \wedge^2 V_\pm$ be the Lie algebras of the holonomy groups of the connections ∇^\pm . Let W be a representation of $\mathfrak{g}_+ \oplus \mathfrak{g}_-$, then d^H sends sections of W into sections of representations that appear in $(\text{Clif}^3(V_+) \oplus \text{Clif}^3(V_-)) \cdot W$.*

Proof. It follows from Lemmata 6.2 and 6.3 that

$$\mathcal{D}_\pm^H : \Gamma(W) \rightarrow \Gamma(\text{Clif}^3(V_\mp) \cdot W),$$

hence, $d^H = \frac{1}{2}(\mathcal{D}_+^H + \mathcal{D}_-^H)$ has the stated property. \square

It follows from the material in Sections 3 and 4 that the difference between an SKT or generalized Kähler structure and parallel (almost) Hermitian or (almost) bi-Hermitian structure, respectively, is that in the former two cases if W is a representation of $\mathfrak{g}_+ \oplus \mathfrak{g}_-$ then $d^H : \Gamma(W) \rightarrow \Gamma(\mathbb{T}_\mathbb{C}M \cdot W)$, while in the latter two cases Proposition 6.4 is the best one can say. This suggests that in general there is a subclass of the space of manifolds (M, g, H) with reduced holonomy which may be of further interest:

Definition 6.5. We say that (∇^+, ∇^-) induces an *integrable* $G_+ \times G_-$ structure if the holonomy of ∇^\pm is G_\pm and

$$d^H : \Gamma(W) \rightarrow \Gamma(\mathbb{T}_\mathbb{C}M \cdot W) \tag{6.5}$$

for every representation $W \subset \wedge^\bullet T_\mathbb{C}^*M$ of $\mathfrak{g}_+ \oplus \mathfrak{g}_-$.

7 Instantons over complex surfaces

Next we use the techniques developed in this section to provide an alternative description of the SKT structure on the moduli space of instantons of a bundle over a complex surface. The tool used to describe this structure are extended actions as introduced in [4] and the SKT reduction theorem as presented in [8]. The argument presented here follows closely the one from [5], so we will spare details and refer to that paper for further reading.

Let $(M, [g], I)$ be a compact complex surface with a conformal Hermitian structure. By a result of Gauduchon [16], there is a representative g of the conformal class which makes (M, g, I) into an SKT manifold, i.e., the corresponding Hermitian form ω satisfies $dd^c\omega = 0$. We let $H = d^c\omega$ and consider $\mathbb{T}M$ endowed with the H -Courant bracket, so that the metric

$$\mathcal{G} = \begin{pmatrix} 0 & g^{-1} \\ g & 0 \end{pmatrix}$$

and the complex structure on V_+ induced by I via the isomorphism $\pi : V_+ \rightarrow TM$ are a positive SKT structure on $\mathbb{T}M$.

Given a bundle E over M with a compact Lie group G as structure group and Lie algebra \mathfrak{g} we let \mathbb{A} be the space of all \mathfrak{g} -connections on E endowed with the trivial 3-form, so that \mathbb{A} is an affine space isomorphic to space of 1-forms on M with values in the adjoint bundle \mathfrak{g}_E , $\Omega^1(M; \mathfrak{g}_E)$. Hence at any connection A we have $T_A\mathbb{A} = \Omega^1(M; \mathfrak{g}_E)$ and, letting κ be a bi-invariant metric on G , we

can use κ to identify $T_A^*\mathbb{A} = \Omega^3(M; \mathfrak{g}_E)$. Indeed, for $X \in \Omega^1(M; \mathfrak{g}_E)$ and $\xi \in \Omega^3(M; \mathfrak{g}_E)$ we define the natural pairing as

$$\xi(X) = 2 \int_M \kappa(X, \xi).$$

Then for $X, Y \in \Omega^1(M; \mathfrak{g}_E)$ and $\xi, \eta \in \Omega^3(M; \mathfrak{g}_E)$ we have

$$\langle X + \xi, Y + \eta \rangle = \int_M \kappa(X, \eta) + \kappa(Y, \xi) = \int_M \kappa(X, \eta) - \kappa(\xi, Y) = \int_M \kappa(X + \xi, Y + \eta)_{Ch}$$

where $\kappa(\cdot, \cdot)_{Ch}$ indicates that one uses κ to pair elements in \mathfrak{g}_E and the Chevalley pairing on forms to obtain a top degree form.

We denote by \mathcal{G} the gauge group of E and by $\tilde{\mathfrak{g}} = \Omega^0(M; \mathfrak{g}_E)$ its Lie algebra. The infinitesimal generator corresponding to $\gamma \in \Omega^0(M; \mathfrak{g}_E)$ at a point $A \in \mathbb{A}$ is just the vector $d_A \gamma \in \Omega^1(M; \mathfrak{g}_E)$ and we can extend this action to form a lifted action as in [8]:

$$\tilde{\Psi} : \Omega^0(M; \mathfrak{g}_E) \longrightarrow \mathbb{T}\mathbb{A} \quad \tilde{\Psi}(\gamma)|_A = d_A^H \gamma = d_A \gamma + H \wedge \gamma,$$

as long as there are no infinitesimal symmetries, i.e., as long as $d_A : \Omega^0(M; \mathfrak{g}_E) \longrightarrow \Omega^1(M; \mathfrak{g}_E)$ has trivial kernel. Therefore, *from this point onwards we only consider connections for which $d_A : \Omega^0(M; \mathfrak{g}_E) \longrightarrow \Omega^1(M; \mathfrak{g}_E)$ has trivial kernel.* If E is a simple $SU(n)$ -bundle then that is the case for all ASD connections.

Next we add a moment map to this action. Our (equivariant) moment map takes values on the \mathcal{G} -module $\mathfrak{h}^* = \Omega_+^2(M; \mathfrak{g}_E)$, the space of self-dual 2-forms:

$$\mu : \mathbb{A} \longrightarrow \Omega_+^2(M; \mathfrak{g}_E) \quad \mu(A) = (F_A)_+,$$

where F_A is the curvature of the connection A and $(\cdot)_+$ indicates projection onto the space of self dual forms, $\Omega_+^2(M; \mathfrak{g}_E)$, which, for 2-forms in 4 dimensions, agrees with the usual self dual 2-forms. If we let \mathfrak{a} be the sum $\tilde{\mathfrak{g}} \oplus \mathfrak{h}$, \mathfrak{a} becomes a Courant algebra if we endow it with the hemisemidirect product:

$$\llbracket (\gamma_1, \lambda_1), (\gamma_2, \lambda_2) \rrbracket = ([\gamma_1, \gamma_2], \gamma_1 \cdot \lambda_2).$$

Then the maps $\tilde{\Psi}$ and μ together give rise to an extended action given infinitesimally by map of Courant algebras:

$$\Psi : \mathfrak{a} \longrightarrow \Gamma(\mathbb{T}\mathbb{A}) \quad \Psi(\gamma, \lambda)|_A = d_A^H \gamma + d\langle \mu, \lambda \rangle.$$

We notice that $\Omega^0(M; \mathfrak{g}_E) + \Omega_+^2(M; \mathfrak{g}_E)$ is isomorphic to $\Omega_+^{ev}(M; \mathfrak{g}_E)$ via the map

$$\gamma + \lambda \mapsto \gamma + \lambda + \star \gamma, \quad \gamma \in \Omega^0(M; \mathfrak{g}_E), \lambda \in \Omega_+^2(M; \mathfrak{g}_E),$$

so we can use $\mathfrak{a} = \Omega_+^{ev}(M; \mathfrak{g}_E)$ and then the extended action is given simply by

$$\Psi : \Omega_+^{ev}(M; \mathfrak{g}_E) \longrightarrow \Omega^{od}(M; \mathfrak{g}_E) \quad \Psi(\alpha)|_A = d_A^H \alpha.$$

Following the reduction procedure, the reduced manifold, \mathcal{M} , is obtained by taking the quotient of $\mu^{-1}(0)$ by the action of the gauge group. In this case, $\mu^{-1}(0)$ consists of the space of anti self-dual connections and hence \mathcal{M} is simply the moduli space of instantons.

The reduction procedure also produces a specific Courant algebroid over the reduced manifold. Namely, if we let \mathbb{K} be the bundle generated by $\Psi(\mathfrak{a})$ and \mathbb{K}^\perp its orthogonal complement with respect to the natural pairing, then the space of \mathcal{G} -invariant sections of $\mathbb{K}^\perp/\mathbb{K}$ over $\mu^{-1}(0)$ inherits a bracket and a nondegenerate pairing which make the quotient bundle $(\mathbb{K}^\perp/\mathbb{K})/\mathcal{G}$ into a Courant algebroid over \mathcal{M} . Notice that at a specific anti self-dual connection A , \mathbb{K} is the image of $\Omega_+^{ev}(M; \mathfrak{g}_E)$ by d_A^H . If we let $d_{A+}^H : \Omega^{od}(M; \mathfrak{g}_E) \longrightarrow \Omega_+^{ev}(M; \mathfrak{g}_E)$ be the composition of d_A^H with projection onto the

SD-forms, then a simple integration by parts shows that $\mathbb{K}^\perp|_A$ is the space of d_{A+}^H -closed forms and hence the reduced Courant algebroid is the degree one cohomology of the following elliptic complex:

$$\{0\} \longrightarrow \Omega_+^{ev}(M; \mathfrak{g}_E) \xrightarrow{d_A^H} \Omega^{od}(M; \mathfrak{g}_E) \xrightarrow{d_{A+}^H} \Omega_+^{ev}(M; \mathfrak{g}_E) \longrightarrow \{0\}. \quad (7.1)$$

In a way, this is the double of the usual elliptic complex describing the tangent space to M :

$$\{0\} \longrightarrow \Omega^0(M; \mathfrak{g}_E) \xrightarrow{d_A} \Omega^1(M; \mathfrak{g}_E) \xrightarrow{d_{A+}} \Omega_+^2(M; \mathfrak{g}_E) \longrightarrow \{0\}. \quad (7.2)$$

Earlier we added the assumption that (7.2) above has no cohomology in degree zero. From now on we also add the assumption that this complex has no cohomology in degree two, so that A corresponds to a smooth point in \mathcal{M} and hence the dimension of \mathcal{M} is given by the index of d_A , which is a topological invariant due to the Atiyah-Singer index theorem. Further, in this case, the cohomology of (7.1) also concentrates in the middle term:

$$H_{d_A^H}^{od}(M; \mathfrak{g}_E) = \frac{\ker(d_{A+}^H : \Omega^{od}(M; \mathfrak{g}_E) \longrightarrow \Omega_+^{ev}(M; \mathfrak{g}_E))}{\text{Im}(d_A^H : \Omega_+^{ev}(M; \mathfrak{g}_E) \longrightarrow \Omega^{od}(M; \mathfrak{g}_E))}.$$

Since M has an SKT structure, we can also endow \mathbb{A} with an SKT structure. Firstly we let \star be the generalized metric: Indeed, in four dimensions $\star^2 = \text{Id}$, spin invariance of the Chevalley pairing means that \star is orthogonal (in even dimensions) and by design we have (c.f. (5.1))

$$\langle \varphi, \star \varphi \rangle = \int_M \kappa(\varphi, \star \varphi)_{Ch} > 0, \quad \text{for } \varphi \in \Omega^{od}(M; \mathfrak{g}_E) \setminus \{0\},$$

so \star is a generalized metric and for this metric V_+ is the space of SD odd forms, $\Omega_+^{od}(M; \mathfrak{g}_E) = (\mathcal{W}^2 + \mathcal{W}^{-2}) \cap \Omega^{od}(M; \mathfrak{g}_E)$, and V_- is the space of ASD odd forms, $\Omega_-^{od}(M; \mathfrak{g}_E) = \mathcal{W}^0 \cap \Omega^{od}(M; \mathfrak{g}_E)$.

For complex structure, we let $\mathbb{I} = e^{\frac{\pi}{2}}$ as in Lemma 1.9. Then that same lemma implies that $\mathbb{I}^2|_{V_+} = -\star = -\text{Id}$. Since this structure is independent of the point $A \in \mathbb{A}$, it is constant and hence integrable. The spaces $V_+^{1,0}$ and $V_+^{0,1}$ can be easily described: since $\mathcal{I} = \mathbb{I}$ on $W^{\pm 2}$ we have $V_+^{1,0} = \mathcal{W}^2 \cap \Omega^{od}(M; \mathfrak{g}_E)$ and $V_+^{0,1} = \mathcal{W}^{-2} \cap \Omega^{od}(M; \mathfrak{g}_E)$.

Further, it is immediate that this SKT structure is invariant by the action of the gauge group. Now, we are in position to use the SKT reduction theorem:

Theorem 7.1 (SKT Reduction Theorem [8]). *Let $\Psi : \mathfrak{a} \longrightarrow \Gamma(\mathbb{T}\mathbb{A})$ be an extended action with moment map μ preserving an SKT structure (\mathbb{G}, \mathbb{I}) on \mathbb{A} as above and let $P = \mu^{-1}(0)$. If the underlying group action on P is free and proper and 0 is a regular value of the moment map, then $\mathcal{M}_{red} = P/\mathcal{G}$ is smooth and the SKT structure on \mathbb{A} reduces to an SKT structure on \mathcal{M}_{red} if and only if $\mathbb{K}^\perp \cap V_+|_P$ is invariant under \mathbb{I} .*

As we observed earlier, at a point $A \in \mathbb{A}$, \mathbb{K}^\perp corresponds to the d_{A+}^H -closed odd forms and hence $\mathbb{K}^\perp \cap V_+$ corresponds to the d_{A+}^H -closed SD odd forms, that is the SD, d_A^H -harmonic odd forms and hence according to the theorem to prove that the moduli space of instantons inherits an SKT from M , we must prove that the space of SD, d_A^H -harmonic odd forms is invariant under \mathbb{I} . To achieve this, one must develop Hodge theory for forms with coefficients. In this case, the condition that the connection is anti self-dual, allows us to achieve the result.

Indeed, similarly to the flat case, we can split d_A^H as a sum of three operators:

$$d_A^H = \delta_+^N + \overline{\delta_+^N} + \not{d}_{A-}^H,$$

$$\delta_+^N : \mathcal{W}^k \longrightarrow \mathcal{W}^{k+2} \quad \overline{\delta_+^N} : \mathcal{W}^k \longrightarrow \mathcal{W}^{k-2} \quad \not{d}_{A-}^H : \mathcal{W}^k \longrightarrow \mathcal{W}^k.$$

Indeed, locally $d_A^H = d^H + A$, for some $A \in \Omega^1(M; \mathfrak{g}_E)$ and hence the splitting of d^H together with the splitting of A into its $V_+^{1,0}$, $V_+^{0,1}$ and V_- components gives the desired decomposition of d_A^H . Just as in Section 5.2, integration by parts gives that $(\delta_+^N)^* = -\overline{\delta_+^N}$ and $\not{d}_{A-}^{H*} = \not{d}_{A-}^H$.

Now, let $\varphi \in (\mathcal{W}^2 \oplus \mathcal{W}^{-2}) \cap \Omega^{od}(M; \mathfrak{g}_E)$. Then the d_A^H -Laplacian computed on φ is given by

$$\begin{aligned} \Delta_H \varphi &= (d_A^{H*} d_{A+}^H + d_A^H d_{A+}^{H*}) \varphi \\ &= (-\delta_+^N - \overline{\delta_+^N} + \not{d}_{A-}^H) \not{d}_{A-}^H \varphi + (\delta_+^N + \overline{\delta_+^N} + \not{d}_{A-}^H) \not{d}_{A-}^H \varphi \\ &= 2(\not{d}_{A-}^H)^2 \varphi. \end{aligned}$$

Hence the Laplacian leaves the spaces $\mathcal{W}^2 \cap \Omega^{od}$ and $\mathcal{W}^{-2} \cap \Omega^{od}$ invariant. Therefore we can decompose the space of harmonic SD odd forms into two spaces $\mathcal{H}^{\pm 2} = \ker(\Delta_H) \cap \mathcal{W}^{\pm 2}$ and \mathbb{I} acts as multiplication by i on \mathcal{W}^2 and by $-i$ on \mathcal{W}^{-2} , so either way it preserves the intersection $\ker(\Delta_H) \cap \mathcal{W}^{\pm 2}$ and hence it preserves the space of SD harmonic odd forms. According to Theorem 7.1, this means that the SKT structure from \mathbb{A} reduces to an SKT structure on \mathcal{M} so we have re-obtained the following result, originally due to Lübke and Teleman [27]:

Theorem 7.2. *Let $(M, I, [g])$ be a compact conformal Hermitian 4-manifold, let E be a bundle over M whose structure group is compact with Lie algebra \mathfrak{g} and let \mathfrak{g}_E be the adjoint bundle over M . Let \mathcal{M}_s be the quotient of the space*

$$\widetilde{P} = \{A \in \mathbb{A} : (F_A)_+ = 0; H_{d_A}^0(M; \mathfrak{g}_E) = H_{d_A}^2(M; \mathfrak{g}_E) = \{0\}\},$$

by the action of the gauge group, i.e., \mathcal{M}_s is the smooth locus of the moduli space of instantons on E . Then \mathcal{M}_s has an SKT structure induced by the unique SKT structure on M in the conformal class $[g]$.

8 Stability

8.1 Stability of SKT structures I

Similarly to the Kähler case, the space of deformations of SKT structures is infinite dimensional even after taking the quotient by the action of the diffeomorphism group. Indeed, from the classical viewpoint, an SKT structure is a Hermitian structure (g, I) for which $dd^c \omega = 0$. For any $f \in C^\infty(M)$ we can consider a new Hermitian form: $\omega + \varepsilon dd^c f$. In a compact manifold if we take ε small enough this gives rise to new non diffeomorphic SKT structures on M still with 3-form $d^c \omega$.

It is therefore more natural to study the deformation problem in the context of stability: Given a generalized complex extension of an SKT structure $(\mathcal{G}, \mathcal{J}_1)$, one is interested in which deformations of \mathcal{J}_1 give rise to deformations of the SKT structure. Since in this setup the SKT structure and the deformation problem are on a fixed Courant algebroid, we are implicitly requiring that the class $[H]$ does not change as the structures vary. The proof of this section's main theorem is inspired on the approach used by Goto to prove the generalized Kähler stability theorem [17].

Before we state the theorem, we must introduce the vector bundles and operators relevant for the deformation theory of SKT manifolds. The bundle that will govern the stability problem is $\wedge^\bullet \overline{L}_2$, where $\overline{L}_2 = V_+^{0,1} \oplus V_-^{1,0}$ is the $-i$ eigenspace of \mathcal{J}_2 . Due to the decomposition of L_2 as a direct sum, we have a bigrading on its exterior algebra, so we define

$$\wedge^{k,l} \overline{L}_2 = \wedge^k V_+^{0,1} \otimes \wedge^l V_-^{1,0}.$$

The relevant differential operators are related to δ_\pm and their complex conjugates, namely we define

$$\begin{aligned} \overline{\partial}_\pm &: \text{Clif}(\mathbb{T}_\mathbb{C}M) \longrightarrow \text{Diff}(\Omega^\bullet(M; \mathbb{C})); & \overline{\partial}_\pm \alpha &= \{\overline{\delta}_\pm, \alpha\}; \\ \partial_\pm &: \text{Clif}(\mathbb{T}_\mathbb{C}M) \longrightarrow \text{Diff}(\Omega^\bullet(M; \mathbb{C})); & \partial_\pm \alpha &= \{\delta_\pm, \alpha\}; \\ \overline{N} &: \text{Clif}(\mathbb{T}_\mathbb{C}M) \longrightarrow \text{Clif}(\mathbb{T}_\mathbb{C}M) & \overline{N} \alpha &= \{\overline{N}, \alpha\}; \end{aligned} \tag{8.1}$$

where $\text{Diff}(\Omega^\bullet(M; \mathbb{C}))$ is the vector space of linear differential operators on forms which is itself a \mathbb{Z}_2 -graded Lie algebra with graded commutator as the bracket.

The main properties of these operators are:

Proposition 8.1. *Let $(\mathcal{G}, \mathcal{J})$ be a generalized Hermitian extension of a positive SKT structure on a manifold with 3-form (M, H) . The restriction of $\bar{\mathcal{N}}$ to $\wedge^\bullet \bar{L}_2$ vanishes and*

$$\bar{\partial}_+ : \Gamma(\wedge^k V_+^{0,1} \otimes \wedge^l V_-^{1,0} \otimes \wedge^m V_-^{0,1}) \longrightarrow \Gamma(\wedge^{k+1} V_+^{0,1} \otimes \wedge^l V_-^{1,0} \otimes \wedge^m V_-^{0,1}); \quad (8.2)$$

$$\partial_- : \Gamma(\wedge^{k,l} \bar{L}_2) \longrightarrow \Gamma(\wedge^{k,l+1} \bar{L}_2). \quad (8.3)$$

Further, the following relations and their complex conjugates hold:

$$\bar{\partial}_+^2 = \bar{\mathcal{N}}^2 = 0; \quad \{\bar{\partial}_+, \bar{\mathcal{N}}\} = 0;$$

$$\begin{aligned} \partial_-^2 = -\{\bar{\mathcal{N}}, \bar{\partial}_+\}; \quad \{\bar{\mathcal{N}}, \partial_-\} = 0; \quad \{\bar{\partial}_-, \bar{\partial}_+\} = 0; \quad \{\bar{\partial}_+, \partial_-\} = -\{\bar{\partial}_-, \bar{\mathcal{N}}\}; \\ \{\partial_+, \bar{\partial}_+\} + \{\partial_-, \bar{\partial}_-\} + \{\mathcal{N}, \bar{\mathcal{N}}\} = 0. \end{aligned}$$

A proof of this proposition is given in the Appendix. Since $\bar{\partial}_+^2 = 0$ we can form the corresponding cohomology:

Definition 8.2. Let $(\mathcal{G}, \mathcal{I})$ be a positive SKT structure on a manifold with 3-form (M, H) and \mathcal{J} be a generalized complex extension of \mathcal{I} . The $\bar{\partial}_+$ -cohomology in \bar{L}_2 , $H_{\bar{\partial}_+}^{\bullet, \bullet}(M; \bar{L}_2)$, is the quotient

$$H_{\bar{\partial}_+}^{k,l}(M; \bar{L}_2) = \frac{\ker(\bar{\partial}_+ : \Gamma(\wedge^{k,l} \bar{L}_2) \longrightarrow \Gamma(\wedge^{k+1,l} \bar{L}_2))}{\text{Im}(\bar{\partial}_+ : \Gamma(\wedge^{k-1,l} \bar{L}_2) \longrightarrow \Gamma(\wedge^{k,l} \bar{L}_2))}$$

Now we are in position to state this section's main theorem:

Theorem 8.3. (SKT Stability Theorem) *Let $(\mathcal{G}, \mathcal{J}_1)$ be a generalized Hermitian extension of a positive SKT structure. Given $\mathcal{J}_{1\varepsilon}$, an analytic family of deformations of \mathcal{J}_1 parametrized by a small disc $D \subset \mathbb{C}$, $\varepsilon : D \longrightarrow \Gamma(\wedge^2 \bar{L}_1)$, we let*

$$\begin{aligned} \varepsilon^+ : D &\longrightarrow \Gamma(\wedge^2 V_+^{0,1}) \\ \varepsilon^\pm : D &\longrightarrow \Gamma(V_+^{0,1} \otimes V_-^{0,1}) \\ \varepsilon^- : D &\longrightarrow \Gamma(\wedge^2 V_-^{0,1}) \end{aligned}$$

be the components of ε and for each of them let $\varepsilon^\bullet = \sum_{k=1}^{\infty} \frac{1}{k!} \varepsilon_k^\bullet t^k$ be the corresponding series expansion. Then the obstructions to finding a generalized metric \mathcal{G}_ε which makes $(\mathcal{G}_\varepsilon, \mathcal{J}_{1\varepsilon})$ into a generalized Hermitian extension of an SKT structure lie in the space $H_{\bar{\partial}_+}^{2,1}(M; \bar{L}_2)$ and the first obstruction is the class $[\partial_- \varepsilon_1^+ + \bar{\mathcal{N}} \varepsilon_1^\pm]$.

Lemma 8.4. There is a map $a : D \subset \wedge^2 \bar{L}_1 \longrightarrow \wedge^2(L_1 \oplus \wedge^2 \bar{L}_1)_{\mathbb{R}}$, where D is a small disc around the origin and $(\cdot)_{\mathbb{R}}$ denotes the real elements in the vector space, such that for each $\varepsilon \in \Gamma(\wedge^2 \bar{L}_1)$, $a(\varepsilon)$ is the unique element in the codomain that satisfies

$$e_*^a \mathcal{J}_1 e_*^{-a} = I_\varepsilon \mathcal{J}_1 I_\varepsilon^{-1}.$$

Further $a_*|_0(\varepsilon) = -(\varepsilon + \bar{\varepsilon})$.

Proof. Indeed, the space of generalized complex structures of the same parity as \mathcal{J}_1 on $T_p M$ is the homogeneous space $SO(\mathbb{T}_p M)/\text{Stab}(\mathcal{J}_1) \cong SO(n, n)/U(n, n)$. Hence, composing the exponential map with the projection

$$\mathfrak{so}(\mathbb{T}_p M) \longrightarrow SO(\mathbb{T}_p M) \longrightarrow SO(\mathbb{T}_p M)/\text{Stab}(\mathcal{J}_1)$$

gives a local submersion. Since the elements in $\mathfrak{so}(\mathbb{T}_p M)$ preserving \mathcal{J}_1 are those in $(L_1 \otimes \overline{L_1})_{\mathbb{R}}$, and $(\wedge^2 L_1 \oplus \wedge^2 \overline{L_1})_{\mathbb{R}}$ is a complementary subspace, we have that

$$(\wedge^2 L_1 \oplus \wedge^2 \overline{L_1})_{\mathbb{R}} \longrightarrow SO(\mathbb{T}_p M) / \text{Stab}(\mathcal{J}_1).$$

is a local diffeomorphism. That is, for each small deformation of a generalized complex structure \mathcal{J}_1 on V there is a unique element $a \in (\wedge^2 L_1 \oplus \wedge^2 \overline{L_1})_{\mathbb{R}}$ which realizes it.

For the last claim, we observe that if $\varepsilon : (-s, s) \longrightarrow \wedge^2 \overline{L_1}$ is a path which passes through 0 at time zero and $v \in L_1$, then $\dot{I}_\varepsilon|_0(v + \bar{v}) = \dot{\varepsilon}|_0(v) + \bar{\varepsilon}|_0(\bar{v})$. On the other hand, for $a : (-s, s) \longrightarrow (\wedge^2 L_1 \oplus \wedge^2 \overline{L_1})_{\mathbb{R}}$, say $a = \alpha + \bar{\alpha}$ with $\alpha \in \wedge^2 L_1$, we have

$$\frac{d}{dt} e_*^a|_0(v + \bar{v}) = \frac{d}{dt} (e^a(v + \bar{v})e^{-a})|_0 = \{a|_0, v + \bar{v}\} = -(\alpha|_0(v) + \bar{\alpha}|_0(\bar{v})),$$

hence if the family a gives rise to the same deformation as ε , we must have $a = -(\varepsilon + \bar{\varepsilon})$. \square

Proof of Theorem 8.3. With this setup, e_*^a is an orthogonal transformation of $\mathbb{T}M$ and hence $e_*^a \mathcal{J}_2 e_*^{-a}$ is an generalized almost complex structure which commutes with $\mathcal{J}_{1\varepsilon}$. If $b \in \mathbb{R} \oplus (L_1 \otimes \overline{L_1})_{\mathbb{R}} \subset \text{Clif}^2(\mathbb{T}M)$ the orthogonal transformation e_*^b preserves \mathcal{J}_1 hence $\mathcal{J}_{1\varepsilon}$ and $\mathcal{J}_{2\varepsilon} = e_*^a e_*^b \mathcal{J}_2 e_*^{-b} e_*^{-a}$ give rise to a generalized Hermitian structure for any choice of b . Our task is to identify suitable conditions on a given a under which we can choose b so that the pair $\mathcal{J}_{1\varepsilon}$ and $\mathcal{J}_{2\varepsilon}$ gives an SKT structure.

Since $\mathcal{J}_{1\varepsilon}$ is integrable, we know that

$$d^H : \mathcal{U}_\varepsilon^{0,n} \longrightarrow \mathcal{U}_\varepsilon^{1,n-1} \oplus \mathcal{U}_\varepsilon^{1,n-3} \oplus \mathcal{U}_\varepsilon^{-1,n-1} \oplus \mathcal{U}_\varepsilon^{-1,n-3}$$

and, according to Theorem 4.9, the SKT condition is that the $\mathcal{U}_\varepsilon^{-1,n-3}$ component of the map above must vanish.

Since $\mathcal{J}_{1\varepsilon}$ and $\mathcal{J}_{2\varepsilon}$ are obtained from \mathcal{J}_1 and \mathcal{J}_2 by the action of $e^a e^b \in \text{Spin}(\mathbb{T}M)$, the corresponding decompositions of spinors are related by

$$\mathcal{U}_\varepsilon^{p,q} = e^a e^b \mathcal{U}^{p,q},$$

so the different components of d^H in the splitting induced by $(\mathcal{J}_{1\varepsilon}, \mathcal{J}_{2\varepsilon})$ (c.f. Figure 3)

$$d^H = \delta_{+\varepsilon} + \bar{\delta}_{+\varepsilon} + \delta_{-\varepsilon} + \bar{\delta}_{-\varepsilon} + N_\varepsilon + \bar{N}_\varepsilon + N_{2\varepsilon} + \bar{N}_{2\varepsilon}$$

are in correspondence with the components of $e^{-b} e^{-a} d^H e^a e^b$ in the splitting determined by $(\mathcal{J}_1, \mathcal{J}_2)$ (see Figure 9).




Figure 9: Nontrivial components of $e^{-b} e^{-a} d^H e^a e^b$ and their relation to the components of d^H with respect to the decomposition of forms determined by $(\mathcal{J}_{1\varepsilon}, \mathcal{J}_{2\varepsilon})$.

Since $a \in (\wedge^2 L_1 \oplus \wedge^2 \overline{L_1})_{\mathbb{R}}$, there are $\alpha^+ \in \Gamma(\wedge^2 V_+^{0,1})$, $\alpha^\pm \in \Gamma(V_+^{0,1} \otimes V_-^{0,1})$ and $\alpha^- \in \Gamma(\wedge^2 V_-^{0,1})$ such that

$$a = \alpha^+ + \alpha^\pm + \alpha^- + \overline{\alpha^+} + \overline{\alpha^\pm} + \overline{\alpha^-}. \quad (8.4)$$

Similarly, any $b \in \Gamma(\mathbb{R} \oplus L_1 \otimes \overline{L_1})_{\mathbb{R}}$ can be decomposed in four components according to the splitting

$$\mathbb{R} \oplus (L_1 \otimes \overline{L_1})_{\mathbb{R}} = \mathbb{R} \oplus (V_+^{1,0} \otimes V_+^{0,1})_{\mathbb{R}} \oplus (V_-^{1,0} \otimes V_-^{0,1})_{\mathbb{R}} \oplus ((V_+^{0,1} \otimes V_-^{1,0}) \oplus (V_+^{1,0} \otimes V_-^{0,1}))_{\mathbb{R}}$$

however, as we will see later, only the last of these components is relevant for the deformation problem. So we let b is of the form

$$b = \beta + \beta^\pm + \overline{\beta^\pm} \quad (8.5)$$

with $\beta^\pm \in \Gamma(V_+^{0,1} \otimes V_-^{1,0})$ and $\beta \in \Gamma(\mathbb{R} \oplus (V_+^{1,0} \otimes V_+^{0,1})_{\mathbb{R}} \oplus (V_-^{1,0} \otimes V_-^{0,1})_{\mathbb{R}})$.

Since the deformation family $\varepsilon(t)$ is analytic on t , so is $a(t)$ and we can try to solve the condition that the $\mathcal{U}^{-1,n-3}$ component of $e^{-b} e^{-a} d^H e^a e^b \psi$ vanishes by a power series argument. In the sequence given an analytic function $f(t)$, where f can be α^+ , α^\pm and so on, we denote the coefficients of its power series expansion by f_k , so that $f = \sum \frac{1}{k!} f_k t^k$.

The first nontrivial condition on a imposed by the integrability of \mathcal{J}_1 is that $[d^H, a_1]$ must have no components in $\mathcal{U}^{n-3, \bullet}$. In terms of the decomposition of a above, this implies

$$[\overline{\delta}_+, \alpha_1^+] = 0 \quad [\overline{\delta}_+, \alpha_1^\pm] + [\overline{\delta}_-, \alpha_1^+] = 0$$

or equivalently

$$\overline{\partial}_+ \alpha_1^+ = 0 \quad \overline{\partial}_+ \alpha_1^\pm + \overline{\partial}_- \alpha_1^+ = 0. \quad (8.6)$$

The first term in the series expansion of $e^{-b} e^{-a} d^H e^a e^b \psi$ is

$$[d^H, b_1 + a_1] \psi.$$

Splitting a and b into their components, we see that the requirement that the $\mathcal{U}^{-1,n-3}$ component vanishes is equivalent to

$$([\overline{\delta}_+, \beta_1^\pm] + [\delta_-, \alpha_1^+] + [\overline{N}, \alpha_1^\pm]) \psi = 0$$

Isolating β^\pm and using that ψ is nonzero, we can rewrite the condition above as

$$\overline{\partial}_+ \beta_1^\pm = -\partial_- \alpha_1^+ - \overline{N} \alpha_1^\pm.$$

And one obvious necessary condition for this to have a solution is that the right hand side must be $\overline{\partial}_+$ -closed. That is indeed the case, since

$$\overline{\partial}_+ (\partial_- \alpha_1^+ + \overline{N} \alpha_1^\pm) = -\partial_- \overline{\partial}_+ \alpha_1^+ - \overline{\partial}_- \overline{N} \alpha_1^+ - \overline{N} \overline{\partial}_- \alpha_1^+ - \overline{N} \overline{\partial}_+ \alpha_1^\pm = 0$$

where for the first equality we used the commutation rule for $\overline{\partial}_+$ and ∂_- from Proposition 8.1 and in the second equality we used equations (8.6) and the fact that $\mathcal{N} \alpha_1^+ = 0$ as $\alpha_1^+ \in \overline{L_2}$. If this $\overline{\partial}_+$ -closed form is in fact $\overline{\partial}_+$ -exact, we can use $G_{\overline{\partial}_+}$, the Green operator for $\overline{\partial}_+$, to find a suitable β_1^\pm :

$$\beta_1^\pm = -\overline{\partial}_+^* G_{\overline{\partial}_+} (\partial_- \alpha_1^+ + \overline{N} \alpha_1^\pm)$$

Finally, we recall that according to Lemma 8.4, $a_1 = -(\varepsilon_1 + \overline{\varepsilon}_1)$, so the components of a_1 are determined by the different components of ε_1 hence the equation β_1^\pm must solve is

$$\overline{\partial}_+ \beta_1^\pm = \partial_- \varepsilon_1^+ + \overline{N} \varepsilon_1^\pm,$$

so the first obstruction class is $[\partial_- \varepsilon_1^+ + \overline{N} \varepsilon_1^\pm] \in H_{\overline{\partial}_+}^{2,1}(M; \overline{L_2})$.

Now we move to the general case, which is proved by induction. We assume that we have chosen a_i and b_i for $i < k$ such that the component of $e^{-b}e^{-a}d^H e^a e^b$ mapping $\mathcal{U}^{0,n}$ into $\mathcal{U}^{-1,n-3}$ vanishes to order $k-1$. Then the vanishing of the order k component of this map is the condition

$$[\beta_k^\pm, \bar{\delta}_+] + [\alpha_k^+, \bar{\delta}_-] + [\alpha_k^\pm, \bar{N}] + F_k(a_1, \dots, a_{k-1}, b_1, \dots, b_{k-1}) = 0,$$

where F_k is some function. This condition is equivalent to

$$\bar{\partial}_+ \beta_k^\pm = -(\partial_- \alpha_k^+ + \bar{N} \alpha_k^\pm + F_k(a_1, \dots, a_{k-1}, b_1, \dots, b_{k-1})).$$

And if there is β_k^\pm which solves this equation, we can proceed to the next step. If for all possible choices of b_i for $i < k$ which guarantee the vanishing of the lower order terms this equation has no solution, then the deformation is obstructed.

Next we show that $\partial_- \alpha_k^+ + \bar{N} \alpha_k^\pm + F_k(a_1, \dots, a_{k-1}, b_1, \dots, b_{k-1})$ is $\bar{\partial}_+$ -closed so we can conclude that the obstruction space is $H^{2,1}(M; \bar{L}_2)$. Since $\bar{\partial}_+ \beta_k^\pm$ is clearly $\bar{\partial}_+$ -closed, we must prove that

$$\bar{\partial}_+(\bar{\partial}_+ \beta_k^\pm + \partial_- \alpha_k^+ + \bar{N} \alpha_k^\pm + F_k(a_1, \dots, a_{k-1}, b_1, \dots, b_{k-1})) = 0$$

but the term in parenthesis is precisely the order k component of $e^{-b}e^{-a}\bar{N}_{2\varepsilon} e^a e^b$ (see Figure 9) and by assumption, the lower order terms of $e^{-b}e^{-a}\bar{N}_{2\varepsilon} e^a e^b$ vanish, therefore we must prove that

$$\bar{\partial}_+ e^{-b}e^{-a}\bar{N}_{2\varepsilon} e^a e^b = \{\bar{\delta}_+, e^{-b}e^{-a}\bar{N}_{2\varepsilon} e^a e^b\} \quad (8.7)$$

vanishes to order k . But since $\bar{\delta}_+ = e^{-b}e^{-a}\bar{\delta}_{+\varepsilon} e^b e^a + O(t)$, we see that (8.7) vanishes to order k if and only if

$$\{e^{-b}e^{-a}\bar{\delta}_{+\varepsilon} e^a e^b, e^{-b}e^{-a}\bar{N}_{2\varepsilon} e^a e^b\}$$

vanishes to order k . But for this operator, we have

$$\{e^{-b}e^{-a}\bar{\delta}_{+\varepsilon} e^a e^b, e^{-b}e^{-a}\bar{N}_{2\varepsilon} e^a e^b\} = e^{-b}e^{-a}\{\bar{\delta}_{+\varepsilon}, \bar{N}_{2\varepsilon}\} e^a e^b = 0,$$

where in the last equality we used that, in a generalized Hermitian manifold, the condition $(d^H)^2 = 0$ gives, among other things, that $\{\bar{\delta}_+, \bar{N}_2\} = 0$.

Finally, if all obstructions vanish, standard elliptic estimates show that the sequence constructed above converges. See for example [17] for the proof of convergence for the analogous problem in the generalized Kähler case. \square

Of course, the same arguments with the obvious changes give deformation results for negative SKT structures. Similarly, Propositions 1.8 and 5.12 can be used to translate our theorem to classical SKT structures:

Theorem 8.5. *Let (M, g, I) be a compact SKT manifold with $d^c \omega = H$. Given I_ε , an analytic family of deformations of the complex structure parametrized by a small disc $D \subset \mathbb{C}$, the obstructions to finding a metric g_ε which makes $(g_\varepsilon, I_\varepsilon)$ into an SKT structure whose 3-form represents the class $[H]$ lie in the Dolbeault cohomology space $H_{\bar{\partial}}^{1,2}(M)$.*

Proof. Under the isomorphism Ψ of Proposition 1.8, \bar{L}_2 is identified with $T_{\mathbb{C}}^* M$. More precisely, we have the following identifications

$$V_+^{0,1} \xrightarrow{\cong} T^{*1,0} M; \quad V_-^{1,0} \xrightarrow{\cong} T^{*0,1} M.$$

Hence $\wedge^{k,l} \bar{L}_2 = \wedge^{k,l} T^* M$ and due to Proposition 5.12, $\bar{\delta}_+$ is identified with ∂ so we have

$$H_{\bar{\partial}_+}^{k,l}(M; \bar{L}_2) = H_{\partial}^{k,l}(M) = H_{\bar{\partial}}^{l,k}(M).$$

In particular we see that the obstruction space is the Dolbeault cohomology $H_{\bar{\partial}}^{1,2}(M)$. \square

Example 8.6. Let M be either a secondary Kodaira, a Hopf or an Inoue surface (the latter with $b_2 = 0$). All these surfaces have the same Betti and Hodge numbers, namely,

$$b_0 = b_1 = 1; \quad b_2 = 0; \quad h^{0,0} = h^{0,1} = 1,$$

and the remaining Betti and Hodge numbers either vanish or are determined by Poincaré and Serre duality. In particular, since $h^{1,2} = 0$ we conclude that SKT structures on these surfaces are stable. Of course, due to Gauduchon's result on the existence of SKT structures on any compact complex surface [16], this conclusion is not particularly strong.

Let N be a product of n such surfaces (with the product complex structure). Due to Künneth's formula for the Dolbeault cohomology, we have that $h^{1,2}(N) = 0$ and hence again the SKT deformation problem is unobstructed and this claim *does not* follow from Gauduchon's theorem. Since we can deform the product complex structure into one that is not a product, this produces nontrivial examples of SKT structures on these manifolds. In particular, we see that $\times_2(S^1 \times S^3)$ with the product complex structure has a stable SKT structure.

Finally, the Dolbeault cohomology of the Lie group $\text{Spin}(4) \cong S^3 \times S^3$ with a left invariant complex structure compatible with a bi-invariant metric is

$$H_{\bar{\partial}}(\text{Spin}(4)) = \wedge \text{span}\{x^{0,1}, z^{2,1}\} \otimes \mathbb{C}[y^{1,1}]/y^2,$$

where $a^{p,q}$ is a generator of bidegree (p, q) [22], hence $H^{1,2}(M)$ is 1-dimensional, generated by $x^{0,1} \cup y^{1,1}$. Using the Künneth formula, we have that $(S^1 \times S^1) \times (S^3 \times S^3)$ with the product structure has $h^{1,2} = 3$, showing that for this structure, the SKT deformation problem is potentially obstructed. The problem is in fact obstructed and one way to understand this is by observing that a left invariant complex structure on $S^3 \times S^3$ compatible with a bi-invariant metric can be deformed into one that is not compatible with any bi-invariant metric and such structure ceases to be SKT. ■

Example 8.7 (Rank two, simple Lie groups). While the proof that a left invariant complex structure on $S^3 \times S^3$ which is not orthogonal with respect to a bi-invariant metric can not be made into an SKT structure seems to be just a computation, for rank two, compact, simple Lie groups the nature of this obstruction is a little clearer.

Firstly, we observe that if I is indeed orthogonal with respect to the bi-invariant metric on a simple Lie group, then the 3-form H is the Cartan 3-form

$$H(X, Y, Z) = \langle [X, Y], Z \rangle.$$

For such a form, the d^H -cohomology of the Lie group vanishes [6].

On the other hand, if G has rank 2, the Dolbeault cohomology of G depends on whether the complex structure is orthogonal with respect to the Killing form or not:

Theorem 8.8 (Pittie [29], Proposition 4.5). *Let G be a rank two, compact, simple Lie group of complex dimension $n + 1$. Then the Dolbeault cohomology of a left invariant complex structure I on G is given, as a graded vector space, by*

$$H_{\bar{\partial}}^{\bullet}(G) = \wedge^{\bullet} \text{span}(x^{0,1}, z^{2,1}) \otimes \mathbb{C}[y^{1,1}]/(y^{1,1})^n$$

if I is compatible with the bi-invariant metric and by

$$H_{\bar{\partial}}^{\bullet}(G) = \wedge^{\bullet} \text{span}(x^{0,1}, z^{n,n-1}) \otimes \mathbb{C}[y^{1,1}]/(y^{1,1})^2$$

otherwise.

In the case when I is compatible with the metric, then $\partial\omega$ represents the class $z^{2,1}$ above and it is immediate that the $\bar{\partial}^{i\partial\omega}$ -cohomology vanishes, as it should, since in an SKT manifold this cohomology is isomorphic to the d^H -cohomology due to Corollary 5.13.

On the other hand, for a complex structure which not compatible with the metric, $H_{\bar{\partial}}^{2,1}(G) = \{0\}$ and hence if the structure were part of an SKT structure then $\partial\omega$ would be $\bar{\partial}$ -exact and the $\bar{\partial}^{i\partial\omega}$ -cohomology would be isomorphic to the Dolbeault cohomology. But since $H_{\bar{\partial}}^{0,1}(G) = \mathbb{C}$ we see that the Dolbeault cohomology is not isomorphic, as a graded vector space, to either the usual de Rham cohomology of G or the d^H cohomology of G . Hence, regardless of the background 3-form chosen, this complex structure is not part of an SKT structure.

Of course, one can deform a complex structure compatible with the metric into one that is not (by changing the choice of complex structure on the Lie algebra of the maximal torus) and the argument above shows that in this case not only do we have $H_{\bar{\partial}}^{1,2}(G) \neq \{0\}$, but the obstruction map is also nontrivial. \blacksquare

8.2 Stability of generalized Kähler structures

Similarly to the SKT deformation problem studied in Section 8.1, a common way to study the question of deformations of a (generalized) Kähler structure $(\mathcal{J}_1, \mathcal{J}_2)$ is in the context of stability: For which deformations of \mathcal{J}_1 is there a corresponding deformation of \mathcal{J}_2 such that the pair of deformed structures is still a (generalized) Kähler structure?

In its classical setting, this question was successfully answered by Kodaira [26], where \mathcal{J}_1 is taken to be the complex structure and in the generalized setting, for analytic deformations, by Goto [17] under the additional hypothesis that the canonical bundle of \mathcal{J}_2 has a globally defined nowhere vanishing closed section. In both cases, there are no obstructions to the problem. However, even in the Kähler setting, if we let \mathcal{J}_1 be the symplectic structure and \mathcal{J}_2 be the complex, the problem does have obstructions [9].

As we have seen in Section 2, a generalized Kähler structure is at the same time a positive and a negative SKT structure. Hence, when studying the stability problem of a generalized Kähler structure one can expect three possible outcomes:

1. The deformation is unobstructed and one can complete the deformed \mathcal{J}_1 to a generalized Kähler structure;
2. The deformation is half obstructed and one can complete the deformed \mathcal{J}_1 to a positive or negative SKT structure;
3. The deformation is fully obstructed and \mathcal{J}_1 can not be completed to either a positive or a negative SKT structure.

Theorem 8.9 (Generalized Kähler Stability Theorem). *Let $(M, \mathcal{J}_1, \mathcal{J}_2)$ be a compact generalized Kähler manifold. Given $\mathcal{J}_{1\varepsilon}$, an analytic family of deformations of \mathcal{J}_1 parametrized by a small disc $D \subset \mathbb{C}$, $\varepsilon : D \rightarrow \Gamma(\wedge^2 \bar{L}_1)$, we let*

$$\begin{aligned} \varepsilon^+ : D &\longrightarrow \Gamma(\wedge^2 V_+^{0,1}) \\ \varepsilon^\pm : D &\longrightarrow \Gamma(V_+^{0,1} \otimes V_-^{0,1}) \\ \varepsilon^- : D &\longrightarrow \Gamma(\wedge^2 V_-^{0,1}) \end{aligned}$$

be the components of ε and for each of them let $\varepsilon^\bullet = \sum_{k=1}^{\infty} \frac{1}{k!} \varepsilon_k^\bullet t^k$ be the corresponding series expansion. Then the obstructions to finding a generalized complex structure $\mathcal{J}_{2\varepsilon}$ which makes $(\mathcal{J}_{1\varepsilon}, \mathcal{J}_{2\varepsilon})$ into a generalized Kähler structure lie in the space

$$\mathcal{H}^{(2,1)+(1,2)} = \frac{\ker(d_{L_2} : \Gamma(\wedge^{2,1} \bar{L}_2 \oplus \wedge^{1,2} \bar{L}_2))}{\text{Im}(d_{L_2} : \Gamma(\wedge^{1,1} \bar{L}_2))},$$

and the first obstruction class is $[\bar{\partial}_+\varepsilon_1^- + \partial_-\varepsilon_1^+]$.

Proof. According to Lemma 8.4, there is an element $a \in \Gamma(\wedge^2 L_1 \oplus \wedge^2 \bar{L}_1)_{\mathbb{R}}$ such that

$$e_*^{-a} \mathcal{J}_1 e_*^a = e_*^{-\varepsilon} \mathcal{J}_1 e_*^{-\varepsilon}$$

and, as in Theorem 8.3, one can compose the orthogonal transformation e_*^a with an orthogonal transformation e_*^b which preserves \mathcal{J}_1 and still obtain the same deformation, that is, for any $b \in \Gamma(L_1 \otimes \bar{L}_1)_{\mathbb{R}}$, $\mathcal{J}_{1\varepsilon}$ and $\mathcal{J}_{2\varepsilon}$ are given by

$$\mathcal{J}_{1\varepsilon} = e_*^{-a} \mathcal{J}_1 e_*^a \quad \text{and} \quad \mathcal{J}_{2\varepsilon} = e^{-a} e^{-b} \mathcal{J}_2 e^b e^a.$$

As before, our quest is to find b for which $\mathcal{J}_{2\varepsilon}$ is integrable and once again we do so using a power series argument. The canonical bundle of $\mathcal{J}_{2\varepsilon}$ is given by $e^a e^b \mathcal{U}^{0,n}$ and integrability is equivalent to

$$e^{-b} e^{-a} d^H e^a e^b : \mathcal{U}^{0,n} \longrightarrow \mathcal{U}^{n-1,1} \oplus \mathcal{U}^{n-1,-1}.$$

If we decompose a and b into their $V_{\pm}^{1,0}$ and $V_{\pm}^{0,1}$ components as in (8.4) and (8.5) and write the corresponding components in power series, then the first condition we must solve is the linear vanishing of the components of $\overline{N_{\mathcal{J}_{2\varepsilon}}}$, the Nijenhuis tensor of $\mathcal{J}_{2\varepsilon}$

$$e^{-b} e^{-a} d^H e^a e^b : \mathcal{U}^{0,n} \longrightarrow \mathcal{U}^{1,n-3} \oplus \mathcal{U}^{-1,n-3}.$$

The linear term which maps $\mathcal{U}^{0,n}$ into $\mathcal{U}^{-1,n-3} \oplus \mathcal{U}^{1,n-3}$ is

$$d_{L_2} \beta_1^{\pm} + \partial_- \alpha_1^+ + \bar{\partial}_+ \bar{\alpha}_1^-$$

So we see that the vanishing of the linear part of the Nijenhuis tensor is equivalent the following condition

$$d_{L_2} \beta_1^{\pm} = -(\bar{\partial}_+ \bar{\alpha}_1^- + \partial_- \alpha_1^+). \quad (8.8)$$

Integrability of $\mathcal{J}_{1\varepsilon}$ means that the $\mathcal{U}^{\pm 3, n-3}$ components of $e^{-a} e^{-b} d^H e^a e^b|_{\mathcal{U}^{0,n}}$ vanish and the linear part of these components is $\bar{\partial}_+ \alpha_1^+ : \mathcal{U}^{0,n} \longrightarrow \mathcal{U}^{-3, n-3}$ and $\partial_- \bar{\alpha}_1^- : \mathcal{U}^{0,n} \longrightarrow \mathcal{U}^{3, n-3}$, hence α_1^+ is $\bar{\partial}_+$ -closed and $\bar{\alpha}_1^-$ is ∂_- -closed, showing that $\bar{\partial}_+ \bar{\alpha}_1^- + \partial_- \alpha_1^+$ is d_{L_2} -closed and hence represents a class in $\mathcal{H}^{(2,1)+(1,2)}$. If this class vanishes, we can find β_1^{\pm} which solves (8.8). That is the first obstruction class is

$$[\bar{\partial}_+ \bar{\alpha}_1^- + \partial_- \alpha_1^+] \in \mathcal{H}^{(2,1)+(1,2)}.$$

Next we assume that we have chosen β_j^{\pm} for $j < k$ such that the Nijenhuis tensor of \mathcal{J}_2 vanishes to order $k-1$ and we must choose β_k^{\pm} which makes the Nijenhuis tensor of \mathcal{J}_2 vanish to order k . Expanding the operator $e^{-b} e^{-a} d^H e^a e^b$ to order k we see that the degree k part of the Nijenhuis tensor is:

$$(\overline{N_{\mathcal{J}_{2\varepsilon}}})_k = d_{L_2} \beta_k^{\pm} + \partial_- \alpha_k^+ + \bar{\partial}_+ \bar{\alpha}_k^- + F_k(a_1, \dots, a_{k-1}, b_1, \dots, b_{k-1}), \quad (8.9)$$

for some function F_k which takes values in $\Gamma(\wedge^{2,1} \bar{L}_2 \oplus \wedge^{1,2} \bar{L}_2)$, and the vanishing of $\overline{N_{\mathcal{J}_{2\varepsilon}}}$ to order k is equivalent to the requirement that the term above vanishes, which can only be achieved if

$$d_{L_2} \beta_k^{\pm} = -(\partial_- \alpha_k^+ + \bar{\partial}_+ \bar{\alpha}_k^- + F_k(a_1, \dots, a_{k-1}, b_1, \dots, b_{k-1})).$$

Next we prove that the right hand side above is d_{L_2} -closed, which allows us to conclude that the obstruction to find β_k^{\pm} lies in $\mathcal{H}^{(2,1)+(1,2)}$.

We let

$$b_{<k+1} = \sum_{j < k+1} \frac{1}{j!} (\beta_j^{\pm} + \bar{\beta}_j^{\mp}) t^j.$$

Since $e^{-a} \mathcal{J}_1 e^a = e^{-a} e^{-b_{<k+1}} \mathcal{J}_1 e^{b_{<k+1}} e^a$ is integrable for any choice of b_k , we have that the operator

$$e^{-b_{<k+1}} e^{-a} d^H e^a e^{b_{<k+1}} |_{\mathcal{U}^{0,n}}$$

has two components, say $d^H |_{\mathcal{U}^{n,0}} = \bar{\partial}_{\mathcal{J}_2 \varepsilon} + \overline{N_{\mathcal{J}_2 \varepsilon}}$,

$$\bar{\partial}_{\mathcal{J}_2 k} = \bar{\partial}_{\mathcal{J}_2} + \bar{v}(t), \quad \bar{v}(t) \in \Gamma(\overline{L_2}), \quad \bar{v}(0) = 0;$$

$$\overline{N_{\mathcal{J}_2 \varepsilon}} = (\overline{N_{\mathcal{J}_2 \varepsilon}})_k t^k + O(t^{k+1}) \in \Gamma(\wedge^{1,2} \overline{L_2}).$$

The condition $(d^H)^2 = 0$ implies $\{\bar{\partial}_{\mathcal{J}_2 \varepsilon}, \overline{N_{\mathcal{J}_2 \varepsilon}}\} = 0$, hence, in particular, the degree k part of this operator vanishes, i.e.,

$$0 = \{\bar{\partial}_{\mathcal{J}_2}, (\overline{N_{\mathcal{J}_2 \varepsilon}})_k\} = d_{L_2}(\overline{N_{\mathcal{J}_2 \varepsilon}})_k.$$

□

Remark. There are natural maps

$$\pi_{2,1} : \mathcal{H}^{(2,1)+(1,2)} \longrightarrow H_{\partial_+}^{2,1}(M; \overline{L_2}) \quad \text{and} \quad \pi_{1,2} : \mathcal{H}^{(2,1)+(1,2)} \longrightarrow H_{\partial_-}^{1,2}(M; \overline{L_2}).$$

given by taking the appropriate components of a class in $\mathcal{H}^{(2,1)+(1,2)}$. It is possible, however, that a nontrivial class in $\mathcal{H}^{(2,1)+(1,2)}$ is mapped into the trivial class by both projections. In terms of deformations, this means that there may be no obstruction to deforming a generalized Kähler structure as either a positive or a negative SKT structure, while the generalized Kähler deformation problem itself is obstructed. A simple example of this phenomenon happens if we regard a Kähler structure (I, ω) as a generalized Kähler structure with \mathcal{J}_1 determined by the symplectic structure and \mathcal{J}_2 by the complex structure: On the one hand not all deformations ω_t of ω can be accompanied by a deformation of I to keep the pair generalized Kähler. On the other hand, keeping I fixed makes the pair (ω_t, I) into a Hermitian symplectic structure as positivity of the tensor $\omega(I, \cdot)$ is an open condition. We can regard this Hermitian symplectic structure as either a positive or a negative SKT structure, hence the obstructions to deformations of the SKT structures vanish.

8.3 Stability of SKT structures II

An inherent part of the setup used for the study of the stability problem of SKT structures in Section 8.1, was that the cohomology class of 3-form was fixed. Depending on the reader's upbringing, that hypothesis may seem unnatural and one might want to study the problem of stability of a classical SKT structure (g, I) without the requirement that $[d^c \omega]$ is fixed. As we will see next, this is actually an easier problem as it has perfectly acceptable solutions which use only classically available tools.

Definition 8.10. Let (M, I) be a complex manifold. The (p, q) -Aeppli cohomology of M is

$$H_A^{p,q}(M) = \frac{\ker(\partial \bar{\partial} : \Omega^{p,q}(M) \longrightarrow \Omega^{p+1,q+1}(M))}{\text{Im}(\partial : \Omega^{p-1,q}(M) \longrightarrow \Omega^{p,q}(M)) + \text{Im}(\bar{\partial} : \Omega^{p,q-1}(M) \longrightarrow \Omega^{p,q}(M))},$$

and the Aeppli numbers are $h_A^{p,q} = \dim(H_A^{p,q}(M))$.

Theorem 8.11. Let (M, g, I) be an SKT structure and for $\varepsilon \in (-\delta, \delta)$, let I_ε be a smooth family of complex structures with $I_0 = I$. If the Aeppli number $h_A^{1,1}$ is constant on ε , then there is a family of metrics g_ε such that $(g_\varepsilon, I_\varepsilon)$ is an SKT structure for small values of ε . In particular, if $h_A^{1,1}$ is constant in a neighbourhood of I in the moduli space of complex structures, then every nearby complex structure is part of an SKT structure.

Proof. If $h_A^{1,1}$ is constant in ε , we can form the family of Green operators G_ε and projection onto harmonics, \mathcal{H}_ε , for the elliptic complex

$$\Omega^{1,0}(M) \oplus \Omega^{0,1} \xrightarrow{\bar{\partial} + \partial} \Omega^{1,1}(M) \xrightarrow{\partial \bar{\partial}} \Omega^{2,2}(M)$$

Since $h_A^{1,1}$ is constant, the form $\omega_\varepsilon = (\mathcal{H}_\varepsilon + (\partial + \bar{\partial})(\partial + \bar{\partial})^* G_\varepsilon) \pi_\varepsilon^{1,1} \omega$ varies smoothly on ε , where $\pi_\varepsilon^{1,1}$ is the projection from $\Omega^2(M)$ onto $\Omega^{1,1}(M)$ with respect to the complex structure I_ε . The operator $(\mathcal{H}_\varepsilon + (\partial + \bar{\partial})(\partial + \bar{\partial})^* G_\varepsilon)$ is just the projection onto $\partial \bar{\partial}$ -closed forms. We see immediately that $\omega_0 = \omega$, that ω_ε is of type $(1,1)$ and $\partial \bar{\partial}$ -closed. Since $\omega(\cdot, I\cdot)$ is positive definite, the same is true for $\omega_\varepsilon(\cdot, I_\varepsilon \cdot)$ as long as I_ε is not too far from I . \square

Example 8.12 (The Iwasawa manifold). The Iwasawa manifold M is the quotient of the 3-dimensional complex Heiseberg group \mathbb{H} by the left action of the subgroup, Λ , of matrices whose entries are Gaussian integers: $M = \mathbb{H}/\Lambda$. The question of existence of SKT structures on the underlying differentiable manifold as well as the stability of such structures was studied by Fino, Parton and Salamon [12]: while the standard complex structure is not part of an SKT structure there are other complex structures on M which are. Further, there are deformations of the SKT complex structures which cease to be SKT.

From our point of view, this is to be expected and amounts to the observation that for the SKT structures, $h_A^{1,1}(M) = 7$ while for generic complex structures $h_A^{1,1}(M) = 6$ [1]. It is also worth noticing that for the families of SKT structures provided by Fino *et al* the cohomology class of the 3-form changes along the family [2]. \blacksquare

9 Appendix - SKT differentials

Throughout this section we let $(M, \mathcal{G}, \mathcal{J})$ be a manifold with a generalized complex extension of an SKT structure and let $\bar{\partial}_\pm, \partial_\pm, \mathcal{N}$ and $\bar{\mathcal{N}}$ be the operators defined in (8.1).

Proposition 9.1. *The restriction of $\bar{\mathcal{N}}$ to $\wedge^\bullet \bar{L}_2$ vanishes and*

$$\bar{\partial}_+ : \Gamma(\wedge^k V_+^{0,1} \otimes \wedge^l V_-^{1,0} \otimes \wedge^m V_-^{0,1}) \longrightarrow \Gamma(\wedge^{k+1} V_+^{0,1} \otimes \wedge^l V_-^{1,0} \otimes \wedge^m V_-^{0,1}); \quad (9.1)$$

$$\partial_- : \Gamma(\wedge^{k,l} \bar{L}_2) \longrightarrow \Gamma(\wedge^{k,l+1} \bar{L}_2). \quad (9.2)$$

Further, the following relations and their complex conjugates hold:

$$\begin{aligned} \bar{\partial}_+^2 &= \bar{\mathcal{N}}^2 = 0; & \{\bar{\partial}_+, \bar{\mathcal{N}}\} &= 0; \\ \partial_-^2 &= -\{\bar{\mathcal{N}}, \bar{\partial}_+\}; & \{\bar{\mathcal{N}}, \partial_-\} &= 0; & \{\bar{\partial}_-, \bar{\partial}_+\} &= 0; & \{\bar{\partial}_+, \partial_-\} &= -\{\bar{\partial}_-, \bar{\mathcal{N}}\}; \\ & & \{\partial_+, \bar{\partial}_+\} &+ \{\partial_-, \bar{\partial}_-\} &+ \{\mathcal{N}, \bar{\mathcal{N}}\} &= 0. \end{aligned}$$

Proof. The relations between the operators follow from the corresponding relations for $\bar{\delta}_+, \delta_-$ and $\bar{\mathcal{N}}$ and the fact that the vector space of operators on forms are a \mathbb{Z}_2 -graded Lie algebra with Lie bracket given by the graded commutator of operators. If we let $\alpha \in \text{Clif}(\mathbb{T}_\mathbb{C}M)$, using the graded Jacobi identity we have

$$\{d^H, \{d^H, \alpha\}\} = \{\{d^H, d^H\}, \alpha\} - \{d^H, \{d^H, \alpha\}\} = -\{d^H, \{d^H, \alpha\}\} = 0,$$

where we have used that $\{d^H, d^H\} = 2(d^H)^2 = 0$. Hence, splitting d^H into its six components, taking $\alpha \in \Gamma(\wedge^j V_+^{1,0} \otimes \wedge^k V_+^{0,1} \otimes \wedge^l V_-^{1,0} \otimes \wedge^m V_-^{0,1})$, applying this operator to elements in $\mathcal{U}^{p,q}$ and imposing the vanishing of the different components of the result we get the relations stated in the theorem.

Since $\bar{\mathcal{N}} \in \Gamma(\wedge^3 \bar{L}_2)$ and \bar{L}_2 is isotropic, $\{\bar{\mathcal{N}}, \alpha\}$ vanishes for every $\alpha \in \Gamma(\wedge^\bullet \bar{L}_2)$.

Next we will prove that (9.1) holds and will omit the proof of (9.2) as it follows the same principles. Firstly, we prove that, when restricted to the domains indicated in (9.1), $\bar{\partial}_+$ actually takes values in the Clifford algebra of $\mathbb{T}M$. Indeed, for $\alpha \in \text{Clif}(\mathbb{T}M)$, $\varphi \in \mathcal{U}^{p,q}$ and $f \in C^\infty(M)$ we decompose df into its $V_+^{1,0}$, $V_+^{0,1}$, $V_-^{1,0}$ and $V_-^{0,1}$ components, say, ∂_+f , $\bar{\partial}_+f$, ∂_-f and $\bar{\partial}_-f$, respectively and compute

$$\begin{aligned}
\{d^H, \alpha\}(f\varphi) &= d^H\alpha f\varphi - (-1)^{|\alpha|}\alpha d^H(f\varphi) \\
&= df \wedge \alpha\varphi + fd^H\alpha \cdot \varphi - (-1)^{|\alpha|}\alpha(df \wedge \varphi + fd^H\varphi) \\
&= (\partial_-f + \bar{\partial}_-f + \partial_+f + \bar{\partial}_+f) \cdot \alpha\varphi + fd^H\alpha\varphi \\
&\quad - (-1)^{|\alpha|}\alpha((\partial_-f + \bar{\partial}_-f + \partial_+f + \bar{\partial}_+f) \cdot \varphi + fd^H\varphi). \\
&= f\{d^H\alpha\}\varphi + (\partial_-f + \bar{\partial}_-f + \partial_+f + \bar{\partial}_+f) \cdot \alpha\varphi \\
&\quad - (-1)^{|\alpha|}\alpha((\partial_-f + \bar{\partial}_-f + \partial_+f + \bar{\partial}_+f) \cdot \varphi).
\end{aligned} \tag{9.3}$$

If $\alpha \in \Gamma(\wedge^k V_+^{0,1} \otimes \wedge^l V_-^{1,0} \otimes \wedge^m V_-^{0,1})$, then $\bar{\partial}_+f$ graded commutes with α and hence if we compose $\{d^H, \alpha\}$ with the projection onto $\mathcal{U}^{p-k+l-m-1, q-k-l+m-1}$ we have

$$\pi_{\mathcal{U}^{p-k+l-m-1, q-k-l+m-1}} \circ \{d^H, \alpha\} : \mathcal{U}^{p,q} \longrightarrow \mathcal{U}^{p-k+l-m-1, q-k-l+m-1}$$

is C^∞ -linear, that is, it is a tensor. Further, using the decomposition d^H into its six components, as in Theorem 4.9, we see that the tensor above corresponds to:

$$\bar{\partial}_+\alpha = \{\bar{\delta}_+, \alpha\} : \mathcal{U}^{p,q} \longrightarrow \mathcal{U}^{p-k+l-m-1, q-k-l+m-1}$$

Since the action of $\text{Clif}(\mathbb{T}M)$ establishes an isomorphism between $\text{Clif}(\mathbb{T}M)$ and the space of endomorphisms of $\wedge^\bullet T^*M$, the tensor $\bar{\partial}_+\alpha$ corresponds to an element in $\text{Clif}(\mathbb{T}M)$:

$$\bar{\partial}_+ : \Gamma(\wedge^k V_+^{0,1} \otimes \wedge^l V_-^{1,0} \otimes \wedge^m V_-^{0,1}) \longrightarrow \text{Clif}(\mathbb{T}_\mathbb{C}M)$$

Next we observe that a Leibniz rule holds. Indeed, not only is the space of graded linear differential operators on $\Omega^\bullet(M)$ a graded Lie algebra with the graded commutator as a bracket, but also this bracket satisfies a Leibniz rule, with respect to the usual composition of operators. In our case, say, for $\alpha, \beta \in \text{Clif}(\mathbb{T}_\mathbb{C}M)$, this translates to

$$\{\bar{\delta}_+, \alpha \circ \beta\} = \{\bar{\delta}_+, \alpha\} \circ \beta + (-1)^{|\alpha|+1}\alpha \circ \{\bar{\delta}_+, \beta\}.$$

Or, in terms of $\bar{\partial}_+$, the Leibniz rule holds:

$$\bar{\partial}_+(\alpha \circ \beta) = \bar{\partial}_+(\alpha) \circ \beta + (-1)^{|\alpha|+1}\alpha \circ \bar{\partial}_+(\beta).$$

Therefore, in order to prove, for example, (9.1) it is enough to establish the result when the argument of $\bar{\partial}_+$ is a section of any of the bundles $V_+^{0,1}$, $V_-^{1,0}$ or $V_-^{0,1}$.

For sections of $\overline{L}_1 = V_+^{0,1} \oplus V_-^{0,1}$, we notice that L_1 is a Lie algebroid whose dual space is naturally isomorphic to \overline{L}_1 hence we have a Lie algebroid differential

$$d_{L_1} : \Gamma(\overline{L}_1) \longrightarrow \Gamma(\wedge^2 \overline{L}_1)$$

and the decomposition $\overline{L}_1 = V_+^{0,1} \oplus V_-^{0,1}$ gives a corresponding decomposition of d_{L_1} in two components. Then one can readily see that $\bar{\partial}_+$ is nothing but the component of d_{L_1} mapping which increases the $V_+^{0,1}$ -degree:

$$\bar{\partial}_+ : \Gamma(V_+^{0,1}) \longrightarrow \Gamma(\wedge^2 V_+^{0,1}), \quad \bar{\partial}_+ : \Gamma(V_+^{0,1}) \longrightarrow \Gamma(V_+^{0,1} \wedge V_-^{0,1}).$$

Therefore to finish the proof we only need to show that

$$\bar{\partial}_+ : \Gamma(V_-^{1,0}) \longrightarrow \Gamma(V_+^{0,1} \otimes V_-^{1,0}). \tag{9.4}$$

Lemma 9.2. The vector bundle $V_-^{1,0}$ is a representation of the Lie algebroid $V_+^{1,0}$ if endowed with the connection

$$\nabla : \Gamma(V_+^{1,0} \otimes V_-^{1,0}) \longrightarrow \Gamma(V_-^{1,0}); \quad \nabla_v w = \pi_{V_-^{1,0}}(\llbracket v, w \rrbracket),$$

where $v \in \Gamma(V_+^{1,0})$ and $w \in \Gamma(V_-^{1,0})$. That is, for $v, v_1, v_2 \in \Gamma(V_+^{1,0})$, $w \in \Gamma(V_-^{1,0})$ and $f \in C^\infty(M)$ we have

$$\begin{aligned} \nabla_{fv} w &= f \nabla_v w, & \nabla_v f w &= (\mathcal{L}_{\pi_T(v)} f) w + f \nabla_v w \\ \nabla_{v_1} \nabla_{v_2} w - \nabla_{v_2} \nabla_{v_1} w - \nabla_{\llbracket v_1, v_2 \rrbracket} w &= 0. \end{aligned}$$

Proof. The first two identities follow from basic properties of the Courant bracket:

$$\nabla_{fv} w = \pi_{V_-^{1,0}}(\llbracket fv, w \rrbracket) = \pi_{V_-^{1,0}}(f \llbracket v, w \rrbracket - (\mathcal{L}_{\pi_T(w)} f) v) = \pi_{V_-^{1,0}}(f \llbracket v, w \rrbracket) = f \nabla_v w.$$

$$\nabla_v f w = \pi_{V_-^{1,0}}(\llbracket v, fw \rrbracket) = \pi_{V_-^{1,0}}(f \llbracket v, w \rrbracket + (\mathcal{L}_{\pi_T(v)} f) w) = f \nabla_v w + (\mathcal{L}_{\pi_T(v)} f) w.$$

Now we prove flatness. Let us first consider the term $\nabla_{v_1} \nabla_{v_2} w$. We decompose $\llbracket v_2, w \rrbracket$ into two components, according to the decomposition $L_1 = V_+^{1,0} \oplus V_-^{1,0}$ and $\nabla_{v_2} w$ is the component in $V_-^{1,0}$:

$$\llbracket v_2, w \rrbracket = u_+^{1,0} + \nabla_{v_2} w, \quad u_+^{1,0} \in \Gamma(V_+^{1,0}).$$

Since $V_+^{1,0}$ is involutive, we see that

$$\llbracket v_1, u_+^{1,0} \rrbracket \in V_+^{1,0}.$$

Hence the $V_-^{1,0}$ component of $\llbracket v_1, \nabla_{v_2} w \rrbracket$ is the $V_-^{1,0}$ component of $\llbracket v_1, \llbracket v_2, w \rrbracket \rrbracket$, that is

$$\nabla_{v_1} \nabla_{v_2} w = \pi_{V_-^{1,0}}(\llbracket v_1, \llbracket v_2, w \rrbracket \rrbracket).$$

Therefore we have

$$\nabla_{v_1} \nabla_{v_2} w - \nabla_{v_2} \nabla_{v_1} w - \nabla_{\llbracket v_1, v_2 \rrbracket} w = \pi_{V_-^{1,0}}(\llbracket v_1, \llbracket v_2, w \rrbracket \rrbracket - \llbracket v_2, \llbracket v_1, w \rrbracket \rrbracket - \llbracket \llbracket v_1, v_2 \rrbracket, w \rrbracket) = 0,$$

where in the last equality we used the fact that the Courant bracket satisfies the Jacobi identity. \square

Since ∇ is a connection, we think of it dually as an operator

$$\nabla : \Gamma(V_-^{1,0}) \longrightarrow \Gamma(V_+^{0,1} \otimes V_-^{1,0}).$$

The following lemma establishes (9.4) and hence finishes the proof of the proposition.

Lemma 9.3. For $\alpha \in \Gamma(V_-^{1,0})$ we have

$$\bar{\partial}_+ \alpha = \nabla \alpha.$$

Proof. Indeed, we check that when paired (i.e., graded commuted) with an element $v \in \Gamma(V_+^{1,0})$ these two operators give the same result. Unwinding the definition of $\bar{\partial}_+$, we have that for $\varphi \in \mathcal{U}^{p,q}$ $\{v, \bar{\partial}_+ \alpha\} \varphi$ is the $\mathcal{U}^{p+1, q-1}$ component of $\{v, \{d^H, \alpha\}\} \varphi$.

$$\begin{aligned} \{v, \bar{\partial}_+ \alpha\} \varphi &= \pi_{U^{p+1, q-1}}(\{v, \{d^H, \alpha\}\} \varphi) \\ &= \pi_{U^{p+1, q-1}}(\llbracket v, \alpha \rrbracket \varphi) \\ &= \pi_{V_-^{1,0}} \llbracket v, \alpha \rrbracket \varphi \\ &= \{v, \nabla \alpha\} \varphi, \end{aligned}$$

where in the second equality we used that the Courant bracket is the derived bracket associated to d^H .

Since $\bar{\partial}_+\alpha$ and $\nabla\alpha$ are both elements in $\text{Clif}(\mathbb{T}_{\mathbb{C}}M)$ we have just established that their difference lies in the annihilator of $V_+^{1,0}$ under graded commutator, that is

$$\nabla\alpha - \bar{\partial}_+\alpha \in \Gamma(\wedge^\bullet V_+^{1,0} \otimes \wedge^\bullet V_-).$$

And we have that when acting in $\mathcal{U}^{p,q}$

$$\nabla\alpha - \bar{\partial}_+\alpha : \mathcal{U}^{p,q} \longrightarrow \mathcal{U}^{p,q-2} \subset \mathcal{W}^{p+q-2}$$

as individually both $\bar{\partial}_+\alpha$ and $\nabla\alpha$ have this property. On the other hand, elements in $\wedge^l V_+^{1,0} \otimes \wedge^m V_-$ map $\mathcal{U}^{p,q} \subset \mathcal{W}^{p+q}$ into \mathcal{W}^{p+q+l} , so the difference $\nabla\alpha - \bar{\partial}_+\alpha$ vanishes and the result follows. \square

References

- [1] D. Angella, *The Cohomologies of the Iwasawa Manifold and of Its Small Deformations*, *J. Geom. Anal.* **23** (2013), 1355–1378.
- [2] A. Baarsma, *Ph.D thesis*, in preparation. Utrecht University.
- [3] J.-M. Bismut, *A local index theorem for non-Kähler manifolds*, *Math. Ann.* **284** (1989), no. 4, 681–699.
- [4] H. Bursztyn, G. R. Cavalcanti, and M. Gualtieri, *Reduction of Courant algebroids and generalized complex structures*, *Adv. Math.* **211** (2007), no. 2, 726–765.
- [5] ———, *Generalized Kähler geometry of instanton moduli space*, 2012. [ArXiv:1203.2385](#).
- [6] G. R. Cavalcanti, *New aspects of the dd^c -lemma*, Ph.D. thesis, Oxford University, 2004.
- [7] ———, *The decomposition of forms and cohomology of generalized complex manifolds*, *J. Geom. Phys.* **57** (2006), no. 1, 121–132.
- [8] ———, *Reduction of metric structures on Courant algebroids*, *J. Symplectic Geom.* **4** (2006), no. 3, 317–343.
- [9] P. de Bartolomeis, *Symplectic deformations of Kähler manifolds*, *J. Symplectic Geom.* **3** (2005), no. 3, 341–355.
- [10] B. de Wit and P. van Nieuwenhuizen, *Rigidly and locally supersymmetric two-dimensional nonlinear σ -models with torsion*, *Nuclear Phys. B* **312** (1989), no. 1, 58–94.
- [11] G. W. Delius, M. Roček, A. Sevrin, and P. van Nieuwenhuizen, *Supersymmetric sigma models with nonvanishing Nijenhuis tensor and their operator product expansion*, *Nuclear Phys. B* **324** (1989), no. 2, 523–531.
- [12] A. Fino, M. Parton, and S. Salamon, *Families of strong KT structures in six dimensions*, *Comment. Math. Helv.* **79** (2004), no. 2, 317–340.
- [13] A. Fino and A. Tomassini, *Blow-ups and resolutions of strong Kähler with torsion metrics*, *Adv. Math.* **221** (2009), no. 3, 914–935.
- [14] A. Frölicher, *Relations between the cohomology groups of Dolbeault and topological invariants*, *Proc. Nat. Acad. Sci. U.S.A.* **41** (1955), 641–644.
- [15] S. J. Gates, Jr., C. M. Hull, and M. Roček, *Twisted multiplets and new supersymmetric nonlinear σ -models*, *Nuclear Phys. B* **248** (1984), no. 1, 157–186.
- [16] P. Gauduchon, *Le théorème de l'excentricité nulle*, *C. R. Acad. Sci. Paris Sér. A-B* **285** (1977), no. 5, A387–A390.
- [17] R. Goto, *Deformations of generalized complex and generalized Kähler structures*, *J. Differential Geom.* **84** (2010), no. 3, 525–560.

- [18] G. Grantcharov, G. Papadopoulos, and Y. S. Poon, *Reduction of HKT-structures*, *J. Math. Phys.* **43** (2002), no. 7, 3766–3782.
- [19] M. Gualtieri, *Generalized geometry and the Hodge decomposition*, 2004. [ArXiv:math/0409093](#).
- [20] ———, *Generalized Kähler geometry*, 2010. [ArXiv:1007.3485](#).
- [21] ———, *Generalized complex geometry*, *Ann. of Math. (2)* **174** (2011), no. 1, 75–123.
- [22] F. Hirzebruch, *Topological methods in algebraic geometry*, Third enlarged edition. New appendix and translation from the second German edition by R. L. E. Schwarzenberger, with an additional section by A. Borel. Die Grundlehren der Mathematischen Wissenschaften, Band 131, Springer-Verlag New York, Inc., New York, 1966.
- [23] N. Hitchin, *Brackets, forms and invariant functionals*, *Asian J. Math.* **10** (2006), no. 3, 541–560.
- [24] C. M. Hull and E. Witten, *Supersymmetric sigma models and the heterotic string*, *Phys. Lett. B* **160** (1985), no. 6, 398–402.
- [25] K. Kodaira, *On the structure of complex analytic surfaces. IV*, *Amer. J. Math.* **90** (1968), 1048–1066.
- [26] K. Kodaira and D. C. Spencer, *On deformations of complex analytic structures. III. Stability theorems for complex structures*, *Ann. of Math. (2)* **71** (1960), 43–76.
- [27] M. Lübke and A. Teleman, *The Kobayashi-Hitchin correspondence*, World Scientific Publishing Co. Inc., River Edge, NJ, 1995.
- [28] A. Neveu and J. H. Schwarz, *Factorizable dual model of pions*, *Nuclear Physics B* **31** (1971), no. 1, 86 – 112.
- [29] H. V. Pittie, *The Dolbeault-cohomology ring of a compact, even-dimensional Lie group*, *Proc. Indian Acad. Sci. Math. Sci.* **98** (1988), no. 2-3, 117–152.
- [30] J. Streets and G. Tian, *A parabolic flow of pluriclosed metrics*, *Int. Math. Res. Not. IMRN* (2010), no. 16, 3101–3133.
- [31] ———, *Regularity theory for pluriclosed flow*, *C. R. Math. Acad. Sci. Paris* **349** (2011), no. 1-2, 1–4.
- [32] C. Voisin, *Hodge theory and complex algebraic geometry. I*, Cambridge Studies in Advanced Mathematics, vol. 76, Cambridge University Press, Cambridge, 2002. Translated from the French original by Leila Schneps.
- [33] J. Wess and B. Zumino, *Consequences of anomalous Ward identities*, *Phys. Lett.* **37B** (1971), 95–97.