



Measure and Integration 2012-13-Selected Solutions Chapter 9

1. **(p.73, exercise 9.1)** Let (X, \mathcal{A}, μ) be a measure space and f a non-negative simple function such that $f(x) = \sum_{j=1}^m y_j 1_{A_j}(x)$, where $y_i \geq 0$ and $A_j \in \mathcal{A}$ are not necessarily disjoint. Show that $I_\mu(f) = \sum_{j=1}^m y_j \mu(A_j)$.

Proof: The function f can be seen as a sum of m simple functions. Hence by Properties 9.3((i) and (ii)), we have

$$I_\mu(f) = I_\mu\left(\sum_{j=1}^m y_j 1_{A_j}\right) = \sum_{j=1}^m I_\mu(y_j 1_{A_j}) = \sum_{j=1}^m y_j I_\mu(1_{A_j}) = \sum_{j=1}^m y_j \mu(A_j).$$

2. **(p.73, exercise 9.3)** Give an example of a sequence (f_n) such that $f_n : \mathbb{R} \rightarrow \mathbb{R}$ is an increasing function, but the sequence (f_n) is not increasing.

Proof: For $k \geq 1$, let $f_k : \mathbb{R} \rightarrow \mathbb{R}$ be given by

$$f_k(x) = \begin{cases} 0 & -\infty < x < -1/k \\ k^3(x + 1/k), & -1/k \leq x \leq 0 \\ k^2, & x \geq 0 \end{cases}$$

Then, it is easy to see that if $x < y$, then $f_k(x) \leq f_k(y)$ so that f_k is an increasing function for each $k \geq 1$. However, the sequence (f_k) is not increasing since for any k , and for any $x \in (-1/k, -1/(k+1))$, one has $f_k(x) > 0$, while $f_{k+1}(x) = 0$ so that $f_{k+1}(x) < f_k(x)$.

3. **(p.73, exercise 9.7)** Let (X, \mathcal{A}) be a measurable space, and $(\mu_j)_{j \in \mathbb{N}}$ a sequence of measures on (X, \mathcal{A}) . Let $\mu = \sum_{j \in \mathbb{N}} \mu_j$ (by problem 4.6, μ is a measure). Show that for every $u \in \mathcal{M}^+(\mathcal{A})$, one has

$$\int u \, d\mu = \sum_{j \in \mathbb{N}} \int u \, d\mu_j.$$

Proof: Suppose first that $u = 1_A$, where $A \in \mathcal{A}$. Then,

$$\int u \, d\mu = \mu(A) = \sum_{n=1}^{\infty} \mu_n(A) = \sum_{n=1}^{\infty} \int u \, d\mu_n.$$

Suppose now that $u = \sum_{k=1}^m a_k 1_{A_k}$ is a non-negative simple function in standard form, note that A_1, \dots, A_m are measurable and disjoint. Then,

$$\int f d\mu = \sum_{k=1}^m a_k \mu(A_k) = \sum_{k=1}^m a_k \sum_{n=1}^{\infty} \mu_n(A_k) = \sum_{n=1}^{\infty} \sum_{k=1}^m a_k \mu_n(A_k) = \sum_{n=1}^{\infty} \int f d\mu_n.$$

Finally, let $u \geq 0$ be measurable. There exists an increasing sequence of non-negative simple functions f_m converging to u . By Theorem 9.6 (Beppo-Levi), $\int u d\mu_j = \lim_{m \rightarrow \infty} \int f_m d\mu_j$ for all $j \in \mathbb{N}$. Consider the double sequence $a_{m,n} = \sum_{j=1}^n \int f_m d\mu_j$. It is easy to see that $(a_{m,n})$ is increasing in m and in n , hence by exercise 4.6, $\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} a_{m,n} = \lim_{n \rightarrow \infty} \lim_{m \rightarrow \infty} a_{m,n}$. Now,

$$\begin{aligned} \int u d\mu &= \lim_{m \rightarrow \infty} \int f_m d\mu \\ &= \lim_{m \rightarrow \infty} \sum_{j=1}^{\infty} \int f_m d\mu_j \\ &= \lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \sum_{j=1}^n \int f_m d\mu_j \\ &= \lim_{n \rightarrow \infty} \lim_{m \rightarrow \infty} \sum_{j=1}^n \int f_m d\mu_j \\ &= \lim_{n \rightarrow \infty} \sum_{j=1}^n \lim_{m \rightarrow \infty} \int f_m d\mu_j \\ &= \lim_{n \rightarrow \infty} \sum_{j=1}^n \int u d\mu_j \\ &= \sum_{j=1}^{\infty} \int u d\mu_j. \end{aligned}$$

4. (**p.73, exercise 9.9**) Let (X, \mathcal{A}, μ) be a measure space, and let $(A_j)_{j \in \mathbb{N}}$ be a sequence of measurable sets. Set

$$\liminf_{j \rightarrow \infty} A_j = \bigcup_{n=1}^{\infty} \bigcap_{j=n}^{\infty} A_j, \quad \text{and} \quad \limsup_{j \rightarrow \infty} A_j = \bigcap_{n=1}^{\infty} \bigcup_{j=n}^{\infty} A_j.$$

- (i) Prove that $\mathbf{1}_{\liminf_{j \rightarrow \infty} A_j} = \liminf_{j \rightarrow \infty} \mathbf{1}_{A_j}$, and $\mathbf{1}_{\limsup_{j \rightarrow \infty} A_j} = \limsup_{j \rightarrow \infty} \mathbf{1}_{A_j}$.
- (ii) Prove that $\mu(\liminf_{j \rightarrow \infty} A_j) \leq \liminf_{j \rightarrow \infty} \mu(A_j)$.
- (iii) Prove that if μ is a finite measure, then $\limsup_{j \rightarrow \infty} \mu(A_j) \leq \mu(\limsup_{j \rightarrow \infty} A_j)$.
- (iv) Provide an example showing that (iii) fails if μ is not finite.

Proof (i): We first begin by proving two general facts, namely if $E = \bigcap_{j=1}^{\infty} E_j$ and

if $F = \bigcup_{j=1}^{\infty} F_j$, then $\mathbf{1}_E = \inf_{n \geq 1} \mathbf{1}_{E_n}$ and $\mathbf{1}_F = \sup_{n \geq 1} \mathbf{1}_{F_n}$. We prove the first, the

second is proved in a similar way. We need to investigate when both sides are equal to 1. To this end, consider

$$\begin{aligned}\mathbf{1}_{\mathbf{E}}(x) = 1 &\iff x \in E_j \text{ for all } j \geq 1 \\ &\iff \mathbf{1}_{E_j}(x) = 1 \text{ for all } j \geq 1 \\ &\iff \inf_{j \geq 1} \mathbf{1}_{E_j}(x) = 1.\end{aligned}$$

This proves that $\mathbf{1}_{\mathbf{E}} = \inf_{n \geq 1} \mathbf{1}_{E_n}$, and similarly $\mathbf{1}_{\mathbf{F}} = \sup_{n \geq 1} \mathbf{1}_{F_n}$. Going back to the proof

of the exercise...we set $B_n = \bigcap_{j=n}^{\infty} A_j$ and $C_n = \bigcup_{j=n}^{\infty} A_j$. Then, $\liminf_{j \rightarrow \infty} A_j = \bigcup_{n=1}^{\infty} B_n$ and

$\limsup_{j \rightarrow \infty} A_j = \bigcap_{n=1}^{\infty} C_n$. By the above we have

$$\mathbf{1}_{\liminf_{j \rightarrow \infty} A_j} = \mathbf{1}_{\bigcup_{n=1}^{\infty} B_n} = \sup_{n \geq 1} \mathbf{1}_{B_n} = \sup_{n \geq 1} \mathbf{1}_{\bigcap_{j=n}^{\infty} A_j} = \sup_{n \geq 1} \inf_{j \geq n} \mathbf{1}_{A_j} = \liminf_{n \rightarrow \infty} \mathbf{1}_{A_j},$$

and

$$\mathbf{1}_{\limsup_{j \rightarrow \infty} A_j} = \mathbf{1}_{\bigcap_{n=1}^{\infty} C_n} = \inf_{n \geq 1} \mathbf{1}_{C_n} = \inf_{n \geq 1} \mathbf{1}_{\bigcup_{j=n}^{\infty} A_j} = \inf_{n \geq 1} \sup_{j \geq n} \mathbf{1}_{A_j} = \limsup_{n \rightarrow \infty} \mathbf{1}_{A_j}.$$

Proof (ii): Applying Fatou's Lemma to the sequence $(\mathbf{1}_{A_j})$ and using part (i), we get

$$\mu(\liminf_{j \rightarrow \infty} A_j) = \int \mathbf{1}_{\liminf_{j \rightarrow \infty} A_j} d\mu = \int \liminf_{j \rightarrow \infty} \mathbf{1}_{A_j} d\mu \leq \liminf_{j \rightarrow \infty} \int \mathbf{1}_{A_j} d\mu = \liminf_{j \rightarrow \infty} \mu(A_j).$$

Proof (iii): Notice that $\mathbf{1}_{A_j} \leq 1$ and $\int 1 d\mu = \mu(X) < \infty$. Hence, by exercise 9.8 (reverse of Fatou's Lemma) we have

$$\limsup_{j \rightarrow \infty} \mu(A_j) = \limsup_{j \rightarrow \infty} \int \mathbf{1}_{A_j} d\mu \leq \int \limsup_{j \rightarrow \infty} \mathbf{1}_{A_j} d\mu = \int \mathbf{1}_{\limsup_{j \rightarrow \infty} A_j} d\mu = \mu(\limsup_{j \rightarrow \infty} A_j).$$

Proof (iv): Consider the measure space $(\mathbb{R}, \mathcal{B}(\mathbb{R}), \lambda)$, where $\mathcal{B}(\mathbb{R})$ is the Borel σ -algebra, and λ is Lebesgue measure. Notice that $\lambda(\mathbb{R}) = \infty$. For $j \geq 1$, let $A_j = [j, 2j]$. Then,

$$\limsup_{j \rightarrow \infty} \mu(A_j) = \inf_{n \rightarrow \infty} \sup_{j \geq n} \mu(A_j) = \inf_{n \rightarrow \infty} \sup_{j \geq n} j = \infty.$$

On the other hand,

$$\mu(\limsup_{j \rightarrow \infty} A_j) = \mu(\bigcap_{n=1}^{\infty} \bigcup_{j=n}^{\infty} [j, 2j]) = \mu(\bigcap_{n=1}^{\infty} [n, \infty)) = \mu(\emptyset) = 0.$$

Hence, (iii) fails in this case.

5. **(p.73, exercise 9.10)** Let (X, \mathcal{A}, μ) be a measure space and (A_j) a sequence of disjoint measurable sets such that $X = \bigcup_{j \in \mathbb{N}} A_j$.

- (i) Show that for every $u \in \mathcal{M}^+(\mathcal{A})$ (i.e. u is a non-negative measurable functions with values in $[0, \infty]$) one has

$$\int u \, d\mu = \sum_{j=1}^{\infty} \int \mathbf{1}_{A_j} u \, d\mu.$$

- (ii) Assume (X, \mathcal{A}, μ) is σ -finite. Use part (i) to construct a measurable function $w > 0$ such that $\int w \, d\mu < \infty$.

Proof (i): From $X = \bigcup_{j \in \mathbb{N}} A_j$ (disjoint union), it is easy to see that $1 = \mathbf{1}_X = \sum_{j=1}^{\infty} \mathbf{1}_{A_j}$. By Corollary 9.9, for any $u \in \mathcal{M}^+(\mathcal{A})$ one has

$$\int u \, d\mu = \int \sum_{j=1}^{\infty} \mathbf{1}_{A_j} u \, d\mu = \sum_{j=1}^{\infty} \int \mathbf{1}_{A_j} u \, d\mu.$$

Proof (ii): Suppose μ is σ -finite, then there exists an increasing sequence (E_n) of measurable sets such that $X = \bigcup_{n \in \mathbb{N}} E_n$, and $\mu(E_n) < \infty$ for all $n \in \mathbb{N}$. Define $A_1 = E_1$, and for $n \geq 2$, $A_n = E_n \setminus E_{n-1}$. Then the sequence (A_n) is disjoint, $X = \bigcup_{j \in \mathbb{N}} A_j$ and $\mu(A_n) = \mu(E_n) - \mu(E_{n-1}) < \infty$. Define w on X by

$$w(x) = \sum_{n=1}^{\infty} \frac{2^{-n}}{\mu(A_n) + 1} \mathbf{1}_{A_n}.$$

Then, clearly, $w(x) > 0$ for all $x \in X$, and by Corollary 9.9,

$$\begin{aligned} \int w \, d\mu &= \int \sum_{n=1}^{\infty} \frac{2^{-n}}{\mu(A_n) + 1} \mathbf{1}_{A_n} \, d\mu = \sum_{n=1}^{\infty} \frac{2^{-n}}{\mu(A_n) + 1} \int \mathbf{1}_{A_n} \, d\mu \\ &= \sum_{n=1}^{\infty} \frac{2^{-n}}{\mu(A_n) + 1} \mu(A_n) \leq \sum_{n=1}^{\infty} 2^{-n} = 1 < \infty. \end{aligned}$$