Conservative Dynamical Systems 2010/2011

Heinz Hanßmann and Holger Waalkens

The last two exercises are homework, to be handed in on Tuesday 9 November.

9.1 Linear Hamiltonian systems

What are the possible types of linear Hamiltonian systems? How do the corresponding Hamiltonian functions look like? Which of the occurring equilibria are structurally stable — and in what sense? Consider in particular the case of one degree of freedom.

9.2 Perturbations of an anharmonic oscillator

Determine the bifurcation diagram of the family

$$H_{\lambda,\mu}(x,y) = \frac{1}{2}y^2 + \frac{1}{24}x^4 + \frac{\lambda}{2}x^2 + \mu x$$

of Hamiltonian systems.

9.3 Symplectic geometry

- 1. Show that a nonzero vector in the symplectic vector space \mathbb{R}^{2n} can be carried into any other nonzero vector by a symplectic transformation.
- 2. Show that not every two-dimensional plane of \mathbb{R}^{2n} can be obtained from a given two-dimensional plane by a symplectic transformation.
- 3. Show that any non-isotropic two-dimensional plane in \mathbb{R}^{2n} can be carried into any other non-isotropic two-dimensional plane by a symplectic transformation.

9.4 Colombo's top

Analyse the dynamics of "Colombo's top" on S^2 , the 2–parameter family with Hamiltonian functions

$$H_{\lambda,\mu}(x,y,z) = -\frac{1}{2}(z-\lambda)^2 + \mu y$$

and Poisson bracket relations $\{x, y\} = z$ plus cyclic permutations.