



## Measure and Integration 2006-Selected Solutions Chapter 9

1. (**p.73, exercise 9.1**) Let  $(X, \mathcal{A}, \mu)$  be a measure space and  $f$  a non-negative simple function such that  $f(x) = \sum_{j=1}^m y_j 1_{A_j}(x)$ , where  $y_i \geq 0$  and  $A_j \in \mathcal{A}$  are not necessarily disjoint. Show that  $I_\mu(f) = \sum_{j=1}^m y_j \mu(A_j)$ .

**Proof:** The function  $f$  can be seen as a sum of  $m$  simple functions. Hence by Properties 9.3((i) and (ii)), we have

$$I_\mu(f) = I_\mu\left(\sum_{j=1}^m y_j 1_{A_j}\right) = \sum_{j=1}^m I_\mu(y_j 1_{A_j}) = \sum_{j=1}^m y_j I_\mu(1_{A_j}) = \sum_{j=1}^m y_j \mu(A_j).$$

2. (**p.73, exercise 9.7**) Let  $(X, \mathcal{A})$  be a measurable space, and  $(\mu_j)_{j \in \mathbb{N}}$  a sequence of measures on  $(X, \mathcal{A})$ . Let  $\mu = \sum_{j \in \mathbb{N}} \mu_j$  (by problem 4.6,  $\mu$  is a measure). Show that for every  $u \in \mathcal{M}^+(\mathcal{A})$ , one has

$$\int u \, d\mu = \sum_{j \in \mathbb{N}} \int u \, d\mu_j.$$

**Proof:** Suppose first that  $u = 1_A$ , where  $A \in \mathcal{A}$ . Then,

$$\int u \, d\mu = \mu(A) = \sum_{n=1}^{\infty} \mu_n(A) = \sum_{n=1}^{\infty} \int u \, d\mu_n.$$

Suppose now that  $u = \sum_{k=1}^m a_k 1_{A_k}$  is a non-negative simple function in standard form, note that  $A_1, \dots, A_m$  are measurable and disjoint. Then,

$$\int f \, d\mu = \sum_{k=1}^m a_k \mu(A_k) = \sum_{k=1}^m a_k \sum_{n=1}^{\infty} \mu_n(A_k) = \sum_{n=1}^{\infty} \sum_{k=1}^m a_k \mu_n(A_k) = \sum_{n=1}^{\infty} \int f \, d\mu_n.$$

Finally, let  $u \geq 0$  be measurable. There exists an increasing sequence of non-negative simple functions  $f_m$  converging to  $f$ . By Theorem 9.6 (Beppo-Levi),  $\int u \, d\mu_j = \lim_{m \rightarrow \infty} \int f_m \, d\mu_j$  for all  $j \in \mathbb{N}$ . Consider the double sequence  $a_{m,n} = \sum_{j=1}^n \int f_m \, d\mu_j$ . It is easy to see that  $(a_{m,n})$  is increasing in  $m$  and in  $n$ , hence by exercise 4.6,  $\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} a_{m,n} = \lim_{n \rightarrow \infty} \lim_{m \rightarrow \infty} a_{m,n}$ . Now,

$$\begin{aligned}
\int u \, d\mu &= \lim_{m \rightarrow \infty} \int f_m \, d\mu \\
&= \lim_{m \rightarrow \infty} \sum_{j=1}^{\infty} \int f_m \, d\mu_j \\
&= \lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \sum_{j=1}^n \int f_m \, d\mu_j \\
&= \lim_{n \rightarrow \infty} \lim_{m \rightarrow \infty} \sum_{j=1}^n \int f_m \, d\mu_j \\
&= \lim_{n \rightarrow \infty} \sum_{j=1}^n \lim_{m \rightarrow \infty} \int f_m \, d\mu_j \\
&= \lim_{n \rightarrow \infty} \sum_{j=1}^n \int u \, d\mu_j \\
&= \sum_{j=1}^{\infty} \int u \, d\mu_j.
\end{aligned}$$

3. (**p.73, exercise 9.9**) Let  $(X, \mathcal{A}, \mu)$  be a measure space, and let  $(A_j)_{j \in \mathbb{N}}$  be a sequence of measurable sets. Set

$$\liminf_{j \rightarrow \infty} A_j = \bigcup_{n=1}^{\infty} \bigcap_{j=n}^{\infty} A_j, \quad \text{and} \quad \limsup_{j \rightarrow \infty} A_j = \bigcap_{n=1}^{\infty} \bigcup_{j=n}^{\infty} A_j.$$

- (i) Prove that  $\mathbf{1}_{\liminf_{j \rightarrow \infty} A_j} = \liminf_{j \rightarrow \infty} \mathbf{1}_{A_j}$ , and  $\mathbf{1}_{\limsup_{j \rightarrow \infty} A_j} = \limsup_{j \rightarrow \infty} \mathbf{1}_{A_j}$ .
- (ii) Prove that  $\mu(\liminf_{j \rightarrow \infty} A_j) \leq \liminf_{j \rightarrow \infty} \mu(A_j)$ .
- (iii) Prove that if  $\mu$  is a finite measure, then  $\limsup_{j \rightarrow \infty} \mu(A_j) \leq \mu(\limsup_{j \rightarrow \infty} A_j)$ .
- (iv) Provide an example showing that (iii) fails if  $\mu$  is not finite.

**Proof (i):** We first begin by proving two general facts, namely if  $E = \bigcap_{j=1}^{\infty} E_j$  and

if  $F = \bigcup_{j=1}^{\infty} F_j$ , then  $\mathbf{1}_E = \inf_{n \geq 1} \mathbf{1}_{E_n}$  and  $\mathbf{1}_F = \sup_{n \geq 1} \mathbf{1}_{F_n}$ . We prove the first, the second is proved in a similar way. We need to investigate when both sides are equal to 1. To this end, consider

$$\begin{aligned}
\mathbf{1}_E(x) = 1 &\iff x \in E_j \text{ for all } j \geq 1 \\
&\iff \mathbf{1}_{E_j}(x) = 1 \text{ for all } j \geq 1 \\
&\iff \inf_{j \geq 1} \mathbf{1}_{E_j}(x) = 1.
\end{aligned}$$

This proves that  $\mathbf{1}_E = \inf_{n \geq 1} \mathbf{1}_{E_n}$ , and similarly  $\mathbf{1}_F = \sup_{n \geq 1} \mathbf{1}_{F_n}$ . Going back to the proof

of the exercise...we set  $B_n = \bigcap_{j=n}^{\infty} A_j$  and  $C_n = \bigcup_{j=n}^{\infty} A_j$ . Then,  $\liminf_{j \rightarrow \infty} A_j = \bigcup_{n=1}^{\infty} B_n$  and

$\limsup_{j \rightarrow \infty} A_j = \bigcap_{n=1}^{\infty} C_n$ . By the above we have

$$\mathbf{1}_{\liminf_{j \rightarrow \infty} A_j} = \mathbf{1}_{\bigcup_{n=1}^{\infty} B_n} = \sup_{n \geq 1} \mathbf{1}_{B_n} = \sup_{n \geq 1} \mathbf{1}_{\bigcap_{j=n}^{\infty} A_j} = \sup_{n \geq 1} \inf_{j \geq n} \mathbf{1}_{A_j} = \liminf_{n \rightarrow \infty} \mathbf{1}_{A_j},$$

and

$$\mathbf{1}_{\limsup_{j \rightarrow \infty} A_j} = \mathbf{1}_{\bigcap_{n=1}^{\infty} C_n} = \inf_{n \geq 1} \mathbf{1}_{C_n} = \inf_{n \geq 1} \mathbf{1}_{\bigcup_{j=n}^{\infty} A_j} = \inf_{n \geq 1} \sup_{j \geq n} \mathbf{1}_{A_j} = \limsup_{n \rightarrow \infty} \mathbf{1}_{A_j}.$$

**Proof (ii):** Applying Fatou's Lemma to the sequence  $(\mathbf{1}_{A_j})$  and using part (i), we get

$$\mu(\liminf_{j \rightarrow \infty} A_j) = \int \mathbf{1}_{\liminf_{j \rightarrow \infty} A_j} d\mu = \int \liminf_{j \rightarrow \infty} \mathbf{1}_{A_j} d\mu \leq \liminf_{j \rightarrow \infty} \int \mathbf{1}_{A_j} d\mu = \liminf_{j \rightarrow \infty} \mu(A_j).$$

**Proof (iii):** Notice that  $\mathbf{1}_{A_j} \leq 1$  and  $\int 1 d\mu = \mu(X) < \infty$ . Hence, by exercise 9.8 (reverse of Fatou's Lemma) we have

$$\limsup_{j \rightarrow \infty} \mu(A_j) = \limsup_{j \rightarrow \infty} \int \mathbf{1}_{A_j} d\mu \leq \int \limsup_{j \rightarrow \infty} \mathbf{1}_{A_j} d\mu = \int \mathbf{1}_{\limsup_{j \rightarrow \infty} A_j} d\mu = \mu(\limsup_{j \rightarrow \infty} A_j).$$

**Proof (iv):** Consider the measure space  $(\mathbb{R}, \mathcal{B}(\mathbb{R}), \lambda)$ , where  $\mathcal{B}(\mathbb{R})$  is the Borel  $\sigma$ -algebra, and  $\lambda$  is Lebesgue measure. Notice that  $\lambda(\mathbb{R}) = \infty$ . For  $j \geq 1$ , let  $A_j = [j, 2j]$ . Then,

$$\limsup_{j \rightarrow \infty} \mu(A_j) = \inf_{n \rightarrow \infty} \sup_{j \geq n} \mu(A_j) = \inf_{n \rightarrow \infty} \sup_{j \geq n} j = \infty.$$

On the other hand,

$$\mu(\limsup_{j \rightarrow \infty} A_j) = \mu(\bigcap_{n=1}^{\infty} \bigcup_{j=n}^{\infty} [j, 2j]) = \mu(\bigcap_{n=1}^{\infty} [n, \infty)) = \mu(\emptyset) = 0.$$

Hence, (iii) fails in this case.

4. (**p.73, exercise 9.10**) Let  $(X, \mathcal{A}, \mu)$  be a measure space and  $(A_j)$  a sequence of disjoint measurable sets such that  $X = \bigcup_{j \in \mathbb{N}} A_j$ .

- (i) Show that for every  $u \in \mathcal{M}^+(\mathcal{A})$  (i.e.  $u$  is a non-negative measurable functions with values in  $[0, \infty]$ ) one has

$$\int u d\mu = \sum_{j=1}^{\infty} \int \mathbf{1}_{A_j} u d\mu.$$

- (ii) Assume  $(X, \mathcal{A}, \mu)$  is  $\sigma$ -finite. Use part (i) to construct a measurable function  $w > 0$  such that  $\int w d\mu < \infty$ .

**Proof (i):** From  $X = \bigcup_{j \in \mathbb{N}} A_j$  (disjoint union), it is easy to see that  $1 = \mathbf{1}_X = \sum_{j=1}^{\infty} \mathbf{1}_{A_j}$ .  
By Corollary 9.9, for any  $u \in \mathcal{M}^+(\mathcal{A})$  one has

$$\int u \, d\mu = \int \sum_{j=1}^{\infty} \mathbf{1}_{A_j} u \, d\mu = \sum_{j=1}^{\infty} \int \mathbf{1}_{A_j} u \, d\mu.$$

**Proof (ii):** Suppose  $\mu$  is  $\sigma$ -finite, then there exists an increasing sequence  $(E_n)$  of measurable sets such that  $X = \bigcup_{n \in \mathbb{N}} E_n$ , and  $\mu(E_n) < \infty$  for all  $n \in \mathbb{N}$ . Define  $A_1 = E_1$ , and for  $n \geq 2$ ,  $A_n = E_n \setminus E_{n-1}$ . Then the sequence  $(A_n)$  is disjoint,  $X = \bigcup_{j \in \mathbb{N}} A_j$  and  $\mu(A_n) = \mu(E_n) - \mu(E_{n-1}) < \infty$ . Define  $w$  on  $X$  by

$$w(x) = \sum_{n=1}^{\infty} \frac{2^{-n}}{\mu(A_n) + 1} \mathbf{1}_{A_n}.$$

Then, clearly,  $w(x) > 0$  for all  $x \in X$ , and by Corollary 9.9,

$$\begin{aligned} \int w \, d\mu &= \int \sum_{n=1}^{\infty} \frac{2^{-n}}{\mu(A_n) + 1} \mathbf{1}_{A_n} \, d\mu = \sum_{n=1}^{\infty} \frac{2^{-n}}{\mu(A_n) + 1} \int \mathbf{1}_{A_n} \, d\mu \\ &= \sum_{n=1}^{\infty} \frac{2^{-n}}{\mu(A_n) + 1} \mu(A_n) \leq \sum_{n=1}^{\infty} 2^{-n} = 1 < \infty. \end{aligned}$$