



## Measure and Integration Solutions 4

1. Let  $A, B \subseteq \mathbb{R}^N$ , and suppose that  $A \subseteq B$  and  $|B \setminus A|_e = 0$ . Show that if  $A$  is measurable, then  $B$  is measurable and  $|A| = |B|$ .

**Proof** Since  $|B \setminus A|_e = 0$ , then  $B \setminus A$  is measurable, and  $B = A \cup (B \setminus A)$  is a union of two measurable sets, hence  $B$  is measurable. Furthermore,  $|A| \leq |B| \leq |A| + |B \setminus A| = |A|$ . Thus,  $|A| = |B|$ .

2. Prove that  $|x + E|_e = |E|_e$  for all  $x \in \mathbb{R}^N$  and every  $E \subseteq \mathbb{R}^N$ .

**Proof** First notice that if  $I = \prod_{k=1}^N [a_k, b_k]$  is a rectangle, and  $x = (x_1, x_2, \dots, x_N) \in \mathbb{R}^N$ , then  $I + x = \prod_{k=1}^N [a_k + x_k, b_k + x_k]$  and  $I - x = \prod_{k=1}^N [a_k - x_k, b_k - x_k]$  are also rectangles, and  $\text{vol}(I) = \text{vol}(I + x) = \text{vol}(I - x)$ . Now, let  $\mathcal{C} = \{I_n\}$  be any countable cover of  $E$  by rectangles, then  $\mathcal{D} = \{I_n + x\}$  is a countable cover of  $E + x$ . Thus,

$$|E + x|_e \leq \sum_{n=1}^{\infty} \text{vol}(I_n + x) = \sum_{n=1}^{\infty} \text{vol}(I_n).$$

Since  $\mathcal{C}$  was an arbitrary cover of  $E$ , it follows that  $|x + E|_e \leq |E|_e$ . The other inequality is proved similarly by starting with a countable cover of  $E + x$  by rectangles.

3. Let  $A \subseteq \mathbb{R}^M$ . The *inner Lebesgue measure* of  $A$  is defined by

$$|A|_i = \sup\{|K|_e : K \subseteq A, K \text{ is compact}\}.$$

Prove the following.

- (a)  $|A|_i \leq |A|_e$  for all  $A \in \mathbb{R}^M$ .
- (b) If  $A \subseteq B$ , then  $|A|_i \leq |B|_i$ .
- (c) If  $A_1, A_2, \dots$  are disjoint, then  $|\bigcup_{n=1}^{\infty} A_n|_i \geq \sum_{n=1}^{\infty} |A_n|_i$ .
- (d) If  $A$  is compact or open, then  $|A|_e = |A|_i$ .

**Proof (a)** Let  $G$  be any open subset containing  $A$ . Then,  $|K|_e \leq |G|_e = |G|$  for any compact subset  $K$  of  $A$ . Hence  $|A|_i \leq |G|_e$ . Since  $G$  is an arbitrary open set containing  $A$ , it follows that  $|A|_i \leq |A|_e$ .

**Proof (b)** Follows from the fact that any compact subset  $K$  of  $A$  is also a compact subset of  $B$ .

**Proof (c)** Fix an integer  $N$ , and let  $K_1, K_2, \dots, K_N$  be any compact subsets of  $A_1, A_2, \dots, A_N$  respectively. Notice that  $K_1, K_2, \dots, K_N$  are pairwise disjoint, and  $\bigcup_{j=1}^N K_j$  is a compact subset of  $\bigcup_{n=1}^{\infty} A_n$ . Then,

$$\sum_{j=1}^N |K_j|_e = \left| \bigcup_{j=1}^N K_j \right|_e \leq \left| \bigcup_{n=1}^{\infty} A_n \right|_i.$$

Taking first the supremum over compact subsets  $K_1 \subseteq A_1$ , then the supremum over compact subsets  $K_2 \subseteq A_2$ , and so on and finally the supremum over compact subsets  $K_N \subseteq A_N$ , one gets

$$\sum_{j=1}^N |A_j|_i \leq \left| \bigcup_{n=1}^{\infty} A_n \right|_i.$$

Finally taking the limit as  $N \rightarrow \infty$ , one has

$$\sum_{j=1}^{\infty} |A_j|_i \leq \left| \bigcup_{n=1}^{\infty} A_n \right|_i.$$

**Proof (d)** Suppose that  $A$  is compact. Since  $A \subseteq A$ , then

$$|A|_e \leq \sup\{|K|_e : K \subseteq A, K \text{ is compact}\} = |A|_i.$$

Combining this with part (a), one gets that  $|A|_e = |A|_i$ .

Suppose that  $A$  is open. By Lemma 2.1.9, there exists a countable non-overlapping exact cover of  $A$  by cubes  $Q_n$ , i.e.,  $A = \bigcup_{n=1}^{\infty} Q_n$ , and hence  $|A|_e \leq \sum_{n=1}^{\infty} |Q_n|_e$ . For any fixed  $N$ ,  $\bigcup_{n=1}^N Q_n$  is a compact subset of  $A$ , hence by Lemma 2.1.1

$$\sum_{n=1}^N |Q_n|_e = \left| \bigcup_{n=1}^N Q_n \right|_e \leq |A|_i.$$

Taking the limit as  $N \rightarrow \infty$ , we have

$$|A|_e \leq \sum_{n=1}^{\infty} |Q_n|_e \leq |A|_i.$$

By part (a), we have  $|A|_e = |A|_i$ .