

METRIC PROPERTIES OF CONFLICT SETS

LEV BIRBRAIR AND DIRK SIERSMA

Communicated by Robert M. Hardt

ABSTRACT. In this paper we show that the tangent cone of a conflict set in \mathbb{R}^n is a linear affine cone over a conflict set of a smaller dimension and has dimension $n - 1$. Moreover we give an example where the conflict sets is not normally embedded and not locally bi-Lipschitz equivalent to the corresponding tangent cone.

1. INTRODUCTION

Let M be a metric space and let $X = \{X_1, \dots, X_k\}$ be a finite collection of closed disjoint nonempty subsets of M . We define a *territory* of a subset $X_i \in X$ with respect to the space M and the collection X in the following way:

$$Ter_M(X_i, X) = \{x \in M \text{ such that } \forall j \quad d(x, X_i) \leq d(x, X_j)\}.$$

We define a *conflict set* of the collection X with respect to M as follows:

$$Conf_M(X) = \{x \in Ter_M(X_i, X) \cap Ter_M(X_j, X) \text{ for some } i \neq j\}.$$

Singularities of conflict sets of collections of disjoint subsets of \mathbb{R}^n is one of natural objects of Singularity Theory. The investigation of conflict sets was initiated by Y. Yomdin [7]. J. Damon [3], P. Giblin and V.M. Zakalyukin [5] and others pointed out that the theory of conflict sets is closely related to other important objects in Singularity Theory: cut loci, medial axes, wave fronts. The results of [3], [6], [8] are devoted to differential geometry of conflict sets.

Here we study properties of general (not necessary generic) singularities of conflict set from a metric viewpoint. Our restriction is so-called "definability". We suppose that all the sets appearing in our investigation are definable in some o-minimal structure [2]. If a reader is not familiar with "o-minimal" language

Research supported under CNPq grant no 300985/93-2.

Research supported under CNPq grant no 420108/2005-8.

he can suppose that all the sets are semialgebraic or subanalytic. If the sets X_i are definable then the same is true for their conflict sets. Thus, all the "good" topological properties hold and tangent cones are well defined.

The main statement of this paper is the structural theorem about the tangent cone of a conflict set. We prove that it is a linear affine cone over a certain conflict set on S^{n-1} (Theorem 2.2). As a corollary of this result we obtain that, for each set X_i of the collection $\{X_1, X_2, \dots, X_k\}$ in \mathbb{R}^2 , the set $Ter X_i$ does not have the cusp-like regions. Moreover, we show that the tangent cone to $Ter X_i$ has a dimension of the ambient space and the tangent cone of $Conf(X)$ has a dimension $n - 1$. A natural question is the following. Is it true or not that conflict sets have "metrically conic" structure near a singular point? More specifically, a conflict set is locally bi-Lipschitz equivalent to the corresponding tangent cone?

The answer is NO: In section 3, we present an example of a collection of sets $\{X_1, X_2\} \subset \mathbb{R}^3$ such that the conflict set of X_1 and X_2 is not locally homeomorphic to its tangent cone and thus not normally embedded in \mathbb{R}^3 .

2. TANGENT CONES OF CONFLICT SETS

Let A be an o-minimal structure over \mathbb{R} . Let X be a collection of definable subsets of a definable set $Y \subset \mathbb{R}^n$ in an o-minimal structure A . Then $Conf_Y(X)$ and $Ter_Y(X_i, X)$ are definable in A sets.

In this paper we are going to suppose that the space \mathbb{R}^n is equipped with the Euclidean metric.

Proposition 2.1. *Let $X = \{X_1, X_2, X_3, \dots, X_k\}$ be a finite collection of closed and definable in an o-minimal structure A subsets of \mathbb{R}^n . Then $\dim Conf_{\mathbb{R}^n}(X) = n - 1$.*

PROOF. Let us first show that if $x_0 \in Conf_{\mathbb{R}^n}(X)$ then there exists a number j such that $x_0 \in \partial Ter_M(X_j, X)$. Let $r_0 = \min d(x_0, X_i)$. Let $\tilde{X} = \{\tilde{X}_1, \tilde{X}_2, \dots, \tilde{X}_k\}$ be a collection of sets (called "supports") defined as follows: $\tilde{X}_i = X_i \cap S_{x_0, r_0}$. Since $x_0 \in Conf_{\mathbb{R}^n}(X)$ we can suppose that there exist two numbers j_1 and j_2 such that \tilde{X}_{j_1} and \tilde{X}_{j_2} are nonempty. Let $x_{j_1} \in \tilde{X}_{j_1}$ and $x_{j_2} \in \tilde{X}_{j_2}$. Then the half-open segment $[x_{j_1}, x_0]$ belongs to $Ter_{\mathbb{R}^n}(X_{j_1}, X)$ and does not belong to $Ter_{\mathbb{R}^n}(X_{j_2}, X)$, the segment (x_{j_2}, x_0) belongs to $Ter_{\mathbb{R}^n}(X_{j_2}, X)$ and does not belong to $Ter_{\mathbb{R}^n}(X_{j_1}, X)$. Hence, x_0 is a boundary point of $Ter_{\mathbb{R}^n}(X_{j_1}, X)$ and of $Ter_{\mathbb{R}^n}(X_{j_2}, X)$. This argument also proves that the sets $Int(Ter_{\mathbb{R}^n}(X_i, X))$ are disjoint. Since the sets $Ter_{\mathbb{R}^n}(X_i, X)$ are definable in the o-minimal structure A , we have: $Conf_{\mathbb{R}^n}(X) \subset \bigcup_i \partial(Ter_{\mathbb{R}^n}(X_i, X))$. That is why $\dim Conf_{\mathbb{R}^n}(X) \leq n - 1$

(see [2]). From the other hand, $\mathbb{R}^n = \bigcup_i \text{Int}(\text{Ter}_{\mathbb{R}^n}(X_i, X)) \bigcup \text{Conf}_{\mathbb{R}^n}(X)$. Since the sets $\text{Int}(\text{Ter}_{\mathbb{R}^n}(X_i, X))$ are disjoint, we obtain that $\dim \text{Conf}_{\mathbb{R}^n}(X) \geq n - 1$. This proves the proposition. \square

A similar statement is true for collections of definable subsets of S^n and is not true for collections of definable subsets of \mathbb{R}^n equipped with New York metric.

Let M be a subset of \mathbb{R}^n . The cone over M with respect to x_0 (notation: $C_{x_0}M$) is a union of all rays connecting x_0 with all the points $y \in M$.

Let $Y \subset \mathbb{R}^n$ be definable in an o-minimal structure A . A tangent cone $T_{x_0}X$ at a point $x_0 \in X$ is the set of all tangent vectors $\frac{d}{dt}\gamma|_{t=0}$ of all definable in A arcs $\gamma: [0, \varepsilon) \rightarrow Y$ such that $\gamma(0) = x_0$ (see [4], [1]). If x_0 is a smooth point of Y then we obtain the definition of the tangent space.

We are going to use another definition of the tangent cone (see also [4]). Let $N_\varepsilon(Y)$ be a set defined as follows:

$$N_\varepsilon(Y) = \frac{1}{\varepsilon} [Y \cap S_{x_0, \varepsilon} - x_0] + x_0.$$

We use the notations: $S_{x_0, \varepsilon}$, for the sphere centered at x_0 with the radius ε ; $B_{x_0, \varepsilon}$, for the closed ball.

Then the following statement is true:

$$T_{x_0}Y = C_{x_0}(\lim_{\varepsilon \rightarrow 0} \text{Hausdorff} N_\varepsilon(Y)).$$

Let $X = \{X_1, \dots, X_k\}$ be a family of closed and disjoint sets on S^{n-1} . Then there is the following relation:

$$\text{Conf}_{\mathbb{R}^n}(X) = C_{o \in \mathbb{R}^n} \text{Conf}_{S^{n-1}}(X).$$

Here we use the standard geodesic metric on S^{n-1} . The proof of this statement is straightforward.

The main result of this paper is the following statement.

Theorem 2.2. *Let $X = \{X_1, \dots, X_k\}$ be a collection of definable in A closed subsets of \mathbb{R}^n such that $X_i \cap X_j = \emptyset$, for $i \neq j$. Let $x_0 \in \text{Conf}_{\mathbb{R}^n}(X)$ and let $r_0 = \min_i d(x_0, X_i)$. Let $\tilde{X} = \{\tilde{X}_1, \tilde{X}_2, \dots, \tilde{X}_k\}$ be a collection of sets (called "the supports") defined as follows: $\tilde{X}_i = X_i \cap S_{x_0, r_0}$. Then the following identities hold:*

- (1) $T_{x_0}(\text{Ter}_{\mathbb{R}^n}(X_i, X)) = C_{x_0}(\text{Ter}_{S_{x_0, r_0}}(\tilde{X}_i, \tilde{X}))$.
- (2) $T_{x_0}(\text{Conf}_{\mathbb{R}^n}(X)) = C_{x_0}(\text{Conf}_{S_{x_0, r_0}}(\tilde{X}))$.

PROOF. Observe that the statement 2 follows immediately from the statement 1 by the definition of conflict sets. Now we are going to show that the germs of $Ter_{\mathbb{R}^n}(X_i, X)$ and of $Conf_{\mathbb{R}^n}(X)$ at x_0 do not change if we cut the sets X_i by balls of the radius bigger than r_0 centered at x_0 . Namely, we prove the following statement:

Lemma 2.3. *Let $X^\varepsilon = \{X_1^\varepsilon, X_2^\varepsilon, \dots, X_k^\varepsilon\}$ be a collection of the sets defined as follows: $X_i^\varepsilon = X_i \cap \bar{B}_{x_0, r_0 + \varepsilon}$. Then, for all i , the germ of $Ter_{\mathbb{R}^n}(X_i, X)$ at x_0 is the same as the germ of the set $Ter_{\mathbb{R}^n}(X_i^\varepsilon, X^\varepsilon)$ at x_0 .*

PROOF. Let z be a point of X_j such that $d(x_0, z) = d(x_0, X_j)$. Take $\delta = \varepsilon/3$. Let $x \in B_{x_0, \delta}$. Let y be a point of X_j such that $d(x, y) = d(x, X_j)$. We are going to show that y actually belong to X_j^ε .

First: $d(x, y) = d(x, X_j) \leq d(x, z) \leq d(x, x_0) + d(x_0, z) \leq \delta + r_0$. Second: suppose that $y \notin X_j^\varepsilon$. Then $d(x, y) > d(x_0, y) - d(x, x_0) = r_0 + \varepsilon - \delta$. This is a contradiction. So, on $B_{x_0, \delta}$ one have: $d(x, X_j) = d(x, X_j^\varepsilon)$ and, therefore, $Ter_{\mathbb{R}^n}(X_i, X) \cap B_{x_0, \delta} = Ter_{\mathbb{R}^n}(X_i^\varepsilon, X^\varepsilon) \cap B_{x_0, \delta}$. \square

We consider "polar coordinates" (ρ, ϕ) near the point x_0 defined as follows. Let $x \in \mathbb{R}^n$ be a point. Set $\phi(x) = \frac{x - x_0}{\|x - x_0\|}$ and $\rho(x) = d(x, x_0)$.

Lemma 2.4 (Shadow lemma). *Let $X = \{X_1, \dots, X_n\}$ be a family of definable sets such that, for all j , $\phi(X_j) = \phi(\tilde{X}_j)$ (we say that X lies in the shadow of the support of X at x_0). Then the germs at x_0 of the sets $Ter_{\mathbb{R}^n}(X_j, X)$ and $Ter_{\mathbb{R}^n}(\tilde{X}_j, \tilde{X})$ are equal, with \tilde{X}_j as in Theorem 2.2.*

PROOF. Let $x_1 \in B_{x_0, r_0/3}$. Then $d(x_1, X_j) = d(x_1, \tilde{X}_j)$. That is why

$$Ter_{\mathbb{R}^n}(\tilde{X}_j, \tilde{X}) \cap B_{x_0, r_0/3} = Ter_{\mathbb{R}^n}(X_j, X) \cap B_{x_0, r_0/3}.$$

\square

Using polar coordinates we define sets Y_i^ε in the following way: $y \in Y_i^\varepsilon$ if, and only if, $y \in B_{x_0, r_0 + \varepsilon}$, $r_0 \leq \rho(y) \leq r_0 + \varepsilon$ and there exists $x \in X_i^\varepsilon$ s.t. $\phi(x) = \phi(y)$. Clearly, $X_i^\varepsilon \subset Y_i^\varepsilon$. Let $Y^\varepsilon = \{Y_1^\varepsilon, Y_2^\varepsilon, \dots, Y_k^\varepsilon\}$. Let $Z^\varepsilon = \{Z_1^\varepsilon, Z_2^\varepsilon, \dots, Z_k^\varepsilon\}$ be a collection of sets defined as follows: $Z_i^\varepsilon = Y_i^\varepsilon \cap S_{x_0, r_0}$. The Hausdorff limits for ε tending to zero of the families Y_i^ε and Z_i^ε are equal to \tilde{X}_i . If $x \in B_{x_0, r_0/3}$ and $d(x, Y_i^\varepsilon) = d(x, y)$, for some $y \in Y_i^\varepsilon$, then $y \in Z_i^\varepsilon$. Clearly, for small ε , we have: $Y_i^\varepsilon \cap Y_j^\varepsilon = \emptyset$, for $i \neq j$, and, thus, $Z_i^\varepsilon \cap Z_j^\varepsilon = \emptyset$. Since all the sets Z_i^ε belong to S_{x_0, r_0} , then $Ter_{\mathbb{R}^n}(Z_i^\varepsilon, Z^\varepsilon) = C_{x_0} Ter_{S_{x_0, r_0}}(Z_i^\varepsilon, Z^\varepsilon)$. Moreover, the germ of the set $Ter_{\mathbb{R}^n}(Y_i^\varepsilon, Y^\varepsilon)$ at the point x_0 is the same as the germ of $Ter_{\mathbb{R}^n}(Z_i^\varepsilon, Z^\varepsilon)$ [Shadow Lemma].

Let us consider the following collection of sets $W^\varepsilon = \{W_1^\varepsilon, W_2^\varepsilon, \dots, W_k^\varepsilon\}$ and $V^\varepsilon = \{V_1^\varepsilon, V_2^\varepsilon, \dots, V_k^\varepsilon\}$ where $W_j^\varepsilon = Y_j^\varepsilon$, for $i \neq j$, and $W_i^\varepsilon = \tilde{X}_i$, $V_j^\varepsilon = \tilde{X}_j$, for $i \neq j$, and $V_i^\varepsilon = Y_i^\varepsilon$. Since $\tilde{X}_s \subset X_s^\varepsilon \subset Y_s^\varepsilon$, for any s , we obtain the following inclusions:

$$\text{Ter}_{\mathbb{R}^n}(X_i^\varepsilon, X^\varepsilon) \subset \text{Ter}_{\mathbb{R}^n}(Y_i^\varepsilon, V^\varepsilon), \quad (1)$$

$$\text{Ter}_{\mathbb{R}^n}(\tilde{X}_i, W^\varepsilon) \subset \text{Ter}_{\mathbb{R}^n}(X_i^\varepsilon, X^\varepsilon).$$

Let \tilde{W}^ε and \tilde{V}^ε be the collections of sets defined as follows: $\tilde{W}^\varepsilon = W^\varepsilon \cap S_{x_0, r_0}$ and $\tilde{V}^\varepsilon = V^\varepsilon \cap S_{x_0, r_0}$. Note, that the germs of the sets $\text{Ter}_{\mathbb{R}^n}(V_j^\varepsilon, V^\varepsilon)$ and $\text{Ter}_{\mathbb{R}^n}(\tilde{V}_j^\varepsilon, \tilde{V}^\varepsilon)$ at x_0 are equal and the germs of $\text{Ter}_{\mathbb{R}^n}(W_j^\varepsilon, W^\varepsilon)$ and $\text{Ter}_{\mathbb{R}^n}(\tilde{W}_j^\varepsilon, \tilde{W}^\varepsilon)$ are also equal (Shadow lemma). The sets $\text{Ter}_{\mathbb{R}^n}(\tilde{V}_i^\varepsilon, \tilde{V}^\varepsilon)$ and $\text{Ter}_{\mathbb{R}^n}(\tilde{W}^\varepsilon, \tilde{W})$ are purely conic, i.e.

$$\text{Ter}_{\mathbb{R}^n}(\tilde{V}_i^\varepsilon, \tilde{V}^\varepsilon) = C_{x_0} \text{Ter}_{S_{x_0, r_0}}(\tilde{V}_i^\varepsilon, \tilde{V}^\varepsilon)$$

and

$$\text{Ter}_{\mathbb{R}^n}(\tilde{W}_i^\varepsilon, \tilde{W}^\varepsilon) = C_{x_0, r_0} \text{Ter}_{S_{x_0, r_0}}(\tilde{W}_i^\varepsilon, \tilde{W}^\varepsilon).$$

Hence,

$$T_{x_0} \text{Ter}_{\mathbb{R}^n}(\tilde{V}_i^\varepsilon, \tilde{V}^\varepsilon) = \text{Ter}_{\mathbb{R}^n}(\tilde{V}_i^\varepsilon, \tilde{V}^\varepsilon)$$

and

$$T_{x_0} \text{Ter}_{\mathbb{R}^n}(\tilde{W}_i^\varepsilon, \tilde{W}^\varepsilon) = \text{Ter}_{\mathbb{R}^n}(\tilde{W}_i^\varepsilon, \tilde{W}^\varepsilon).$$

Using the inclusions (1) we obtain:

$$T_{x_0}(\text{Ter}_{\mathbb{R}^n}(X_i, X)) = T_{x_0}(\text{Ter}_{\mathbb{R}^n}(X_i^\varepsilon, X^\varepsilon)) \subset \text{Ter}_{\mathbb{R}^n}(\tilde{V}_i^\varepsilon, \tilde{V}^\varepsilon), \quad (2)$$

$$\text{Ter}_{\mathbb{R}^n}(\tilde{W}_i^\varepsilon, \tilde{W}^\varepsilon) \subset T_{x_0}(\text{Ter}_{\mathbb{R}^n}(X_i^\varepsilon, X^\varepsilon)) = T_{x_0}(\text{Ter}_{\mathbb{R}^n}(X_i, X)).$$

Taking the Hausdorff limit in the inclusions (2) we obtain

$$T_{x_0}(\text{Ter}_{\mathbb{R}^n}(X_i, X)) \subset C_{x_0}(\text{Ter}_{S_{x_0, r_0}}(\tilde{X}_i, \tilde{X}))$$

and

$$C_{x_0}(\text{Ter}_{S_{x_0, r_0}}(\tilde{X}_i, \tilde{X})) \subset T_{x_0}(\text{Ter}_{\mathbb{R}^n}(X_i, X)).$$

This proves the theorem. \square

Proposition 2.5. *Let $X = \{X_1, \dots, X_k\}$ be a collection of definable in A sets in \mathbb{R}^n . Then*

- (1) $T_{x_0}(\text{Ter}_{\mathbb{R}^n}(X_i, X))$ has a nonempty interior, for all i such that $x_0 \in \text{Ter}(X_i, X)$.
- (2) If $x_0 \in \text{Conf}_{\mathbb{R}^n}(X)$ then $\dim(T_{x_0} \text{Conf}_{\mathbb{R}^n}(X)) = n - 1$.

1. If $\tilde{X}_i \neq \emptyset$ then $Ter_{S^{n-1}}(\tilde{X}_i, \tilde{X})$ has a nonempty interior. Thus, by Theorem 2.2, $T_{x_0}(Ter_{\mathbb{R}^n}(X_i, X))$ has a nonempty interior. [2] By Proposition 2.1 (see also the remark), $\dim Conf_{S^{n-1}}(\tilde{X}) = n-2$. Hence, $\dim(T_{x_0}Conf_{\mathbb{R}^n}(X)) = n-1$. \square

Theorem 2.6 ("no cusp" property in \mathbb{R}^2). *Let $X = \{X_1, \dots, X_k\}$ be a collection of definable in A sets on \mathbb{R}^2 . Let $Y = Conf_{\mathbb{R}^2}(X)$. Let $y_0 \in Y$. Then the germ of Y at y_0 is a collection of definable in A differentiable arcs $\{\gamma_1, \gamma_2, \dots, \gamma_s\}$ such that y_0 belongs to each γ_i and the unit tangent vectors of γ_i and γ_j are different, for $i \neq j$.*

PROOF. Let $\gamma_1: [0, \varepsilon) \rightarrow Conf_{\mathbb{R}^2}(X)$ be a definable nonconstant arc such that $\gamma_1(0) = y_0$ and $|\gamma_1(t) - y_0| = t$. Since the sets X_i are definable in the o-minimal structure A we can find two sets $X_1, X_2 \in X$ such that $\gamma_1 \subset Conf_{\mathbb{R}^2}(\bar{X})$ where $\bar{X} = \{X_1, X_2\}$. Let l be the tangent ray to γ_1 at y_0 . By Theorem 2.2, l is a bisector ray of the angle defined by points $x_1 \in \tilde{X}_1$, $x_2 \in \tilde{X}_2$ and y_0 , where x_1, x_2 are boundary points on S_{y_0, r_0} of the supporting sets \tilde{X}_1 and \tilde{X}_2 on S_{y_0, r_0} .

Let $\delta > 0$ be a sufficiently small number such that the sets X_1^δ and X_2^δ - their radial projections to the supporting circle are disjoint. Let $\gamma_2: [0, \varepsilon) \rightarrow Conf_{\mathbb{R}^2}(\bar{X})$ be another definable in A arc such that $|\gamma_2(t) - y_0| = t$ and the germs at y_0 of the sets $\Gamma_1 = \gamma_1([0, \varepsilon))$ and $\Gamma_2 = \gamma_2([0, \varepsilon))$ are different and Γ_2 is also tangent to l at y_0 . By Arc Selection Lemma (see [2]), there exist two pairs of definable in A arcs $\alpha_1, \alpha_2: [0, \varepsilon) \rightarrow X_1^\delta$ and $\beta_1, \beta_2: [0, \varepsilon) \rightarrow X_2^\delta$ such that $d(\gamma_1(t), X_1) = |\gamma_1(t) - \alpha_1(t)|$, $d(\gamma_2(t), X_1) = |\gamma_2(t) - \alpha_2(t)|$, $d(\gamma_1(t), X_2) = |\gamma_1(t) - \beta_1(t)|$ and $d(\gamma_2(t), X_2) = |\gamma_2(t) - \beta_2(t)|$. If $\alpha_1(t) = \alpha_2(t) = x_1$ and $\beta_1(t) = \beta_2(t) = x_2$, then the germs of γ_1 and γ_2 at y_0 are equal to the germ of l at y_0 . Thus, we can suppose that $\alpha_1(t) \neq Const$, for small t , and that, for small $t \neq 0$ we have $d(\gamma_1(t), X_1) > d(\gamma_2(t), X_1)$.

Take $t > 0$ sufficiently small. The segment connecting $\gamma_1(t)$ and $\beta_1(t)$ intersects the arc $\gamma_2(t)$. Let z be an intersection point. Observe that $\beta_1(t)$ realizes the shortest distance between z and X_1 . Since t is small and β_1 is a definable in A arc, then we can suppose that $\beta_1(t)$ is a smooth point.

Consider now the circle $S_{\gamma_1(t), |\gamma_1(t) - \beta_1(t)|}$ and the circle $S_{z, |z - \beta_1(t)|}$. These circles are tangent at the point $\beta_1(t)$. That is why the ball with the center at z and the radius $|z - \beta_1(t)|$ does not contain any point of X_1 . But it means that z does not belong to $Conf_{\mathbb{R}^2}(\bar{X})$. It is a contradiction. \square

3. AN EXAMPLE OF NOT NORMALLY EMBEDDED CONFLICT SET

Here we are going to construct an example of a family of sets $X_1, X_2 \in \mathbb{R}^3$ satisfying the following conditions:

- a) $Conf_{\mathbb{R}^3}(\{X_1, X_2\})$ is not normally embedded in \mathbb{R}^3 .
- b) There exists a point $x_0 \in Conf_{\mathbb{R}^3}(\{X_1, X_2\})$ such that, for small r , we have that $B_{x_0, r} \cap Conf_{\mathbb{R}^3}(\{X_1, X_2\})$ is not homeomorphic to $T_{x_0}(Conf_{\mathbb{R}^3}(\{X_1, X_2\}))$.

Example 3.1. Consider the space \mathbb{R}^3 with coordinates (x_1, x_2, x_3) . Let $X_1 \subset \mathbb{R}^3$ be a union of the hyperplanes: $x_3 = 1$ and $x_3 = -1$. Let X_2 be a union of the points: $a_1 = (1, 0, 0)$ and $a_2 = (-1, 0, 0)$.

Theorem 3.2. The set $Conf_{\mathbb{R}^3}(\{X_1, X_2\})$ satisfies the conditions a) and b) described above.

PROOF. The set $Conf_{\mathbb{R}^3}(\{X_1, X_2\})$ can be obtained as follows. Let $Y_1 \subset \mathbb{R}^2$ be the conflict set of the point $a_1 = (1, 0)$ and the union of straight lines $x_3 = 1$ and $x_3 = -1$. Observe, that here we consider \mathbb{R}^2 with coordinates (x_1, x_3) . The set Y_1 is a union of a part of the parabola defined by the point a_1 and the line $x_3 = 1$ situated above the line $x_3 = 0$ and a part of the parabola defined by the same point a_1 and the line $x_3 = -1$ situated below the line $x_3 = 0$.

Let $Y_2 \subset \mathbb{R}^2$ be the conflict set of the point $a_2 = (-1, 0)$ and the union of the lines $x_3 = 1$ and $x_3 = -1$. The set Y_2 can be obtained from Y_1 by the transformation: $(x_1, x_3) \rightarrow (-x_1, x_3)$. The set $Conf_{\mathbb{R}^3}(\{X_1, X_2\})$ can be obtained as a union of the revolution surface of Y_1 with respect to the straight line $x_1 = 1, x_2 = 0$ and the revolution surface of Y_2 with respect to the line $x_1 = -1, x_2 = 0$. The intersection of the set $Conf_{\mathbb{R}^3}(\{X_1, X_2\})$ with the plane $x_3 = 0$ is a union of two metric copies of S^1 . These circles are tangent at the origin. That is why the set $Conf_{\mathbb{R}^3}(\{X_1, X_2\})$ is not normally embedded. The tangent cone $T_0 Conf_{\mathbb{R}^3}(\{X_1, X_2\})$ is a union of two planes intersecting transversally. The germ of the set $Conf_{\mathbb{R}^3}(\{X_1, X_2\})$ is homeomorphic to the quotient space of the disjoint union of two copies of \mathbb{R}^2 by the identification of the two origins. \square

Acknowledgements. The first author was supported by CNPq grant N 300985/93-2, the second author was supported by CNPq grant N 420108/2005-8.

We would like to thank Y. Yomdin, T. Mostowski, A. Fernandes and M. van Manen for their interest on this work.

REFERENCES

- [1] Birbrair, L. "Local bi-Lipschitz classification of 2-dimensional semialgebraic sets." - Houston Journal of Math., v.25, N3 (1999), 453-472.
- [2] Van den Dries, L. "Tame topology and o-minimal structures." - London Mathematical Society Lecture Note Series, 248. Cambridge University Press, Cambridge (1998), 180pp.

- [3] Damon, J. "Smoothness and geometry of boundaries associated to skeletal structures II." - Geometry in the Blum case. *Compos. Math.* 140 (2004), no. 6, 1657–1674.
- [4] Federer, H. "Geometric measure theory." - Die Grundlehren der mathematischen Wissenschaften, Band 153 Springer-Verlag New York Inc., New York 1969, 676 pp.
- [5] Giblin, P. J.; Zakalyukin, V. M. "Singularities of center symmetry sets." - *Proc. London Math. Soc.* (3) 90 (2005).
- [6] Siersma, D. "Properties of conflict sets in the plane." - Geometry and topology of caustics—CAUSTICS '98 (Warsaw), 267–276, Banach Center Publ., 50, Polish Acad. Sci., Warsaw, 1999.
- [7] Yomdin, Y. "On the local structure of a generic central set." - *Compositio Math.* 43 (1981), no. 2, 225–238.
- [8] van Manen, M. "The Geometry of conflict Sets. Dissertation"; Universiteit Utrecht 2003; 111p. Ph.D. Thesis.

Received April 17, 2007

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DO CEARÁ (UFC), CAMPUS DO PICICI, BLOCO 914, CEP. 60455-760. FORTALEZA-CE, BRASIL

E-mail address: birb@ufc.br

DEPARTMENT OF MATHEMATICS, UTRECHT UNIVERSITY, P.O.Box 80.010, 3508 TA UTRECHT, THE NETHERLANDS

E-mail address: siersma@math.uu.nl