Take home exercise 3, alternative formulation

Let S^1 be the unit circle in the complex plane. For $w = w(\varphi) = e^{i\varphi} \in S^1$ we define the following rotations about the *z*-axis and about the *y*-axis in \mathbb{R}^3 ,

$$R_w := \begin{pmatrix} \cos \varphi & -\sin \varphi & 0 \\ \sin \varphi & \cos \varphi & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

and

$$r_w := \begin{pmatrix} \cos(\varphi + \frac{\pi}{2}) & 0 & -\sin(\varphi + \frac{\pi}{2}) \\ 0 & 1 & 0 \\ \sin(\varphi + \frac{\pi}{2}) & 0 & \cos(\varphi + \frac{\pi}{2}) \end{pmatrix} = \begin{pmatrix} -\sin\varphi & 0 & -\cos\varphi \\ 0 & 1 & 0 \\ \cos\varphi & 0 & -\sin\varphi \end{pmatrix}.$$

Let

$$X = S^1 \times [-1, 1],$$

equipped with the restriction of the Euclidean topology on $\mathbb{C} \times \mathbb{R}$. We define the map $f: S^1 \times [-1,1] \to \mathbb{R}^3$ by

$$f(w,t) = R_{w^2}(2e_1 + r_w(te_3)),$$
 (**)

where e_j denotes the j-th standard basis vector in \mathbb{R}^3 . Here the index w^2 is on purpose; note that it equals $w(2\varphi)$.

- (a) Argue that the image M of f is a geometric realization of the Möbius band in \mathbb{R}^3 . See also Exercise 1.12.
- (b) Determine the equivalence relation R on X which turns $f: X \to M$ into a quotient modulo R. Determine an action of the group $\mathbb{Z}_2 = \{1, -1\}$ on X whose orbits are precisely the equivalence classes of R.
- (c) Show that there exists a continuous bijection $F: X/\mathbb{Z}_2 \to M$. (Later we will see that by compactness of X this implies that F is a homeomorphism).
- (d) We consider the continuous map

$$h: [0,1] \times [-1,1] \to X = S^1 \times [-1,1], \quad h(s,t) = (e^{i\pi s},t).$$

Let $p: X \to X/\mathbb{Z}_2$ be the natural projection. Show that $p \circ h$ is a continuous surjection from $[0,1] \times [-1,1]$ onto X/\mathbb{Z}^2 .

(e) Describe the gluing relation G on $[0,1] \times [-1,1]$ for which $p \circ h$ is a quotient modulo G. Show that $([0,1] \times [-1,1])/G$ is homeomorphic to X/\mathbb{Z}_2 .